

**Title:**

**Production of Se-enriched *Chlorella* biomass grown in heterotrophic and phototrophic regimes**

**Authors & affiliations:**

**Mykola Mylenko**<sup>1</sup>, Jan Kuta<sup>2</sup>, Dai Long Vu<sup>1</sup>, Karolína Ranglová<sup>1</sup>, David Kubáč<sup>1</sup>, Martin Pablo Caporgno<sup>1</sup>, Jiří Kopecký<sup>1,3</sup>, Jiří Masojídek<sup>1</sup> and Pavel Hrouzek<sup>1</sup>

<sup>1</sup>Centre Algatech, Institute of Microbiology, Třeboň, Czech Republic;

<sup>2</sup>Research Centre for Toxic Compounds in the Environment (RECETOX), Masaryk University, Brno, Czech Republic

<sup>3</sup>Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic

**Abstract:**

Selenium (Se) is an essential micro-nutrient that is heterogeneously distributed in nature [1, 2]. In areas with low environmental levels of Se, nutritional supplementation of this trace element can prevent its deficiency and improve specific physical and physiological conditions in humans and domestic animals [2, 3].

Microalgal biomass enriched with organically bound Se in feedstuffs, represents a better source of this element for humans and animals relative to potentially toxic, inorganic Se-salts [2, 3].

In this work, the relationship between physiological changes and selenoaminoacid accumulation were studied in *Chlorella* cultures grown in the presence of Se in heterotrophic (strain G-120) or phototrophic (strain R-117; CCALA 1107) regimes. LC-ICP-MS was used to quantify selenomethionine, selenocysteine and methylselenocysteine, while ICP-MS was used to assess Se content. It was found that cultivation medium containing Se with a concentration of 25 mg L<sup>-1</sup> did not have any deleterious effect on microalgae growth, but it induced a high level of selenoaminoacid accumulation. Based on these results we developed the chemostat-based process for continuous production of Se-enriched *Chlorella* biomass in a fermentor (heterotrophic regime). Obtained kinetic data can be used to efficiently upscale the cultivation for industrial needs. The chemostat-based process was also compared with the large-scale production of Se-enriched biomass in outdoor, thin-layer cascades (phototrophic regime). In conclusion, the results reveal the potential use of this technology for the production of Se-enriched biomass as a food and feed supplement.

**References**

- [1] Ullah H. et. al. (2018) Ecotoxicol. Environ. Saf. 149, 291–306.
- [2] Doucha J. et. al. (2009) Appl. Microbiol. Biotechnol. 83, 1001–1008.
- [3] Hosnedlova B. et. al. (2017) Int. J. Mol. Sci. 18, 2209.

**Acknowledgements**

This research was supported by the Ministry of Education of the Czech Republic, project Algatech Plus LO1416.