

Recent Advances in the Detection of Food Toxins Using Mass Spectrometry

Published as part of *Chemical Research in Toxicology* virtual special issue "Mass Spectrometry Advances for Environmental and Human Health".

Vishal Ahuja, Amanpreet Singh, Debarati Paul, Diptarka Dasgupta, Petra Urajová, Sounak Ghosh, Roshani Singh, Gobardhan Sahoo, Daniela Ewe, and Kumar Saurav*

Cite This: *Chem. Res. Toxicol.* 2023, 36, 1834–1863


Read Online

ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: Edibles are the only source of nutrients and energy for humans. However, ingredients of edibles have undergone many physicochemical changes during preparation and storage. Aging, hydrolysis, oxidation, and rancidity are some of the major changes that not only change the native flavor, texture, and taste of food but also destroy the nutritive value and jeopardize public health. The major reasons for the production of harmful metabolites, chemicals, and toxins are poor processing, inappropriate storage, and microbial spoilage, which are lethal to consumers. In addition, the emergence of new pollutants has intensified the need for advanced and rapid food analysis techniques to detect such toxins. The issue with the detection of toxins in food samples is the nonvolatile nature and absence of detectable chromophores; hence, normal conventional techniques need additional derivatization. Mass spectrometry (MS) offers high sensitivity, selectivity, and capability to handle complex mixtures, making it an ideal analytical technique for the identification and quantification of food toxins. Recent technological advancements, such as high-resolution MS and tandem mass spectrometry (MS/MS), have significantly improved sensitivity, enabling the detection of food toxins at ultralow levels. Moreover, the emergence of ambient ionization techniques has facilitated rapid in situ analysis of samples with lower time and resources. Despite numerous advantages, the widespread adoption of MS in routine food safety monitoring faces certain challenges such as instrument cost, complexity, data analysis, and standardization of methods. Nevertheless, the continuous advancements in MS-technology and its integration with complementary techniques hold promising prospects for revolutionizing food safety monitoring. This review discusses the application of MS in detecting various food toxins including mycotoxins, marine biotoxins, and plant-derived toxins. It also explores the implementation of untargeted approaches, such as metabolomics and proteomics, for the discovery of novel and emerging food toxins, enhancing our understanding of potential hazards in the food supply chain.



CONTENTS

1. Introduction	1835	4. Recent Advancements in Food Toxin Detection Using MS	1839
2. Conventional Methods for the Detection of Food Toxins	1836	4.1. Mycotoxins	1839
2.1. Immunoassays	1836	4.2. Bacterial Toxins	1843
2.2. Chromatographic Techniques	1837	4.2.1. Detection of Bacterial Food Toxins	1844
2.3. Spectroscopic-Based Techniques: UV–visible and Fluorescence Spectroscopy	1837	4.3. Marine Biotoxins	1845
2.4. Biological Assays	1837		
2.5. Biochemical Assays	1837		
2.6. Microbiological Assays	1837		
2.7. Sensor-Based Approaches	1837		
3. Mass Spectrometry-Based Detection	1837		

Received: August 25, 2023
Revised: October 30, 2023
Accepted: November 8, 2023
Published: December 7, 2023



Table 1. Various Food Toxins from Various Sources

Class	Toxin's name	Source	Effect	Ref
Mycotoxins	Aflatoxin	<i>Aspergillus flavus</i> and <i>A. parasiticus</i>	Liver failure, cirrhosis	16
	Lysergic acid (ergot alkaloids)	<i>Claviceps purpurea</i>	Ergotism, vasoconstriction, uterine contraction	17
	Fumonisin B1 and B2	<i>Fusarium verticillioides</i> and <i>Fusarium proliferatum</i>	disruption of sphingolipid metabolism, leuko-encephalomalacia	18
	Ochratoxin A	<i>Aspergillus</i> and <i>Penicillium</i>	Carcinogenic, immunotoxic mutagenic, nephrotoxic, and teratogenic	19
	Patulin	<i>Aspergillus</i> , <i>Byssoschlamys</i> and <i>Penicillium</i>	Teratogenic, carcinogenic and mutagenic	20
	Zearalenone	<i>Fusarium graminearum</i> , <i>F. culmorum</i> , <i>F. crookwellense</i> , <i>F. poae</i> , <i>F. semitectum</i> , and <i>F. equiseti</i>	Hepatotoxicity, immunotoxicity, reproductive toxicity	21
	Tentoxin	<i>Alternaria</i>	Genotoxic, mutagenic, and carcinogenic	22
Bacterial toxins	Cholera toxins	<i>Vibrio cholerae</i>	diarrhea	23
	Enterotoxins	<i>Staphylococcus epidermidis</i>	Toxic shock syndrome	24
	Shiga toxins	<i>Escherichia coli</i>	Gastrointestinal complications	25
	Botulinum toxins	<i>Clostridium botulinum</i>	Neurotoxic	26
	Cereulide	<i>Bacillus cereus</i>	Dysfunction of liver, pancreatic islet, intestines, brain,	27
Marine biotoxins	Saxitoxin	Cyanobacteria and dinoflagellates	Neurotoxin, paralysis	28
	domoic acid	Diatoms	Neurotoxin	29
	Azaspiracid	<i>Azadiniumpoporum</i>	Diarrheic shellfish poisoning	30
	Brevetoxin	<i>Karenia brevis</i>	Immunotoxicity	31
	okadaic acid	<i>Halichondriamelanodocia</i> and <i>Halichondriaokadae</i>	Diarrhea, nausea	32
Plant-based toxins	Cyanogenic glycosides	Almonds, cassava, pome fruit, stone fruit	Tissue damage	33
	Furocoumarins	Citrus fruits	Skin cancer	34
	Ptaquiloside	Bracken ferns	Carcinogenic	35
	Dehydropyrrolizidine	<i>Cyanoglossum</i> , <i>Senecio</i> , <i>Echium</i> , <i>Crotalaria</i> , <i>Heliotropium</i> , <i>Symphytum</i> , <i>Trichodesma</i>	Carcinogenic	36

4.3.1. MS Analysis and Detection of Algal and Marine Biotoxins	1847
4.4. Phytotoxins	1849
4.5. Emerging Toxins	1849
5. Detection of Food Fraud and Food Adulteration	1851
6. Current Challenges and Future Prospects	1851
6.1. Representative Sample	1851
6.2. Sample Preparation	1851
6.3. Low Concentration	1852
6.4. Complex Matrix	1852
6.5. Portability of Instrument	1852
6.6. Cost	1852
6.7. Multiple Toxins Contamination	1852
7. Future Perspective	1853
Author Information	1853
Corresponding Author	1853
Authors	1853
Author Contributions	1853
Notes	1853
Biographies	1854
References	1856

1. INTRODUCTION

Serious health hazards and outbreaks resulting from food spoilage are major concerns of food safety worldwide.¹ Due to significant biological activity and poor detection by conventional testing techniques, food toxins among other contaminants constitute a serious risk to the public's health.^{2,3} Food toxins are compounds that can contaminate different food products during manufacturing, processing, transit, or storage.⁴ According to Fletcher and Netzel, these poisons can come from a variety of sources including fungi, bacteria, algae, plants, and animals.⁵ Mycotoxins produced by molds, marine biotoxins from toxic

algal blooms, and plant-derived toxins like alkaloids and glycoalkaloids are all well-known examples of food toxins.^{6,7} Mycotoxins are mainly produced by toxigenic fungal species belonging to the genera of *Fusarium*, *Aspergillus*, and *Penicillium*.⁸ These mycotoxins pose a challenge to food safety because they can contaminate food products even when good storage and processing protocols are employed for food safety. Mycotoxin contamination accounts for the major cause of food borne diseases as reported by the World Health Organization (WHO). Recent studies performed by the European Commission revealed that 80% of the samples were contaminated with at least one mycotoxin.⁹ Perusal of literature revealed that most of the mycotoxins are chemically and thermally stable; therefore, they can survive under storage, processing, and even cooking.¹⁰ It has been observed that crops that are stored for more than a few days become probable targets for the growth of fungi and mycotoxin formation. These mycotoxins can affect a variety of food commodities such as dried fruits, coffee, spices, nuts, cereals, oil seeds, fruits, spices, cocoa, beans, etc.¹¹ Apart from mycotoxins, other microorganisms like bacteria, algae, and even plants also produce such metabolites that are equally toxic and lethal. Bacteria contamination is mainly attributed to poor hygiene and cleanliness and common uncleaned and poorly cooked meat and vegetable-based foods, especially fermented foods. Nem Chua fermented food prepared from pork sausage in Vietnam has the possibility for *Staphylococcus aureus* contamination, and New Zealand mussel (*Perna canaliculus*) traditional fermented food of New Zealand is usually contaminated with *Clostridium botulinum*.¹² Likewise, aquatic and marine food like Shellfish, fish, and even water are contaminated with algal biotoxins.¹³ In the case of plants, some normal metabolites produced for various purposes like natural defense and stress tolerance act as toxins for other organisms. Cyanogenic glycosides in almonds, and summer fruits, furocoumarins in




Figure 1. Schematic representation of various methods used for the detection of food toxins.

citrus fruits, and lectins in beans are some of the common examples of phytotoxins.¹⁴ Besides, some newly emerging chemicals include perchlorate, flame retardants, halo compounds, packaging materials, petrochemicals residues, health-care products traces, and microplastics.¹⁵ The use of contaminated cereals, grapes, and barley used to produce wine and beer products and their consumption are the main causes of toxicological effects in human beings (Table 1).

The use of contaminated cereals, grapes, and barley for the production of wine and beer products is the cause of toxicological effects in human beings. Consumption of contaminated meat and milk-based products is another route for these toxins to enter the human food chain. Further, the abusive use of drugs in livestock, animal waste pollution, and the use of industrial wastewater for irrigation are also responsible factors for the entry of these contaminants into the food chain.³⁷ Conventional methods for the detection of food toxin include immunoassays and chromatographic techniques, which, while effective, have certain limitations.³⁸ Immunoassays, for instance, can be sensitive but may give false results if structurally related compounds are present in the testing matrix.¹⁰ Chromatographic techniques, on the other hand, require complex sample preparation and longer analysis times.³⁹ In contrast, significant advancements in analytical techniques have revolutionized the field of food safety, and one such breakthrough is the application of mass spectrometry (MS) for the rapid and sensitive detection of food toxins. It is highly sensitive and provides selectivity and capability to handle complex mixtures, making it an ideal tool for the detection and characterization of food toxins.^{40,41} This led to the selection of the MS-based approach as the first choice tool among researchers, regulatory agencies, and food industries to ensure the safety and quality of the food supply chain.^{41–43}

Recent advances and innovations in instrumentation, such as the development of high-resolution mass spectrometry (HRMS) and tandem mass spectrometry (MS/MS), have significantly improved sensitivity and selectivity, allowing for the

detection of food toxins at ultralow levels.⁴⁴ Additionally, several ionization methods allow rapid *in situ* analysis, reducing the time and resources required for analysis.^{45,46} Further, advancement in the field of untargeted metabolomics and several web-based repositories of metabolites allows for the detection of not only the known toxin but also the unknown variants of toxins, broadening our understanding of potential hazards in the food supply chain.^{47,48} The current review summarizes the overview of available detection techniques of food toxins and then further elaborates on the MS-based approaches, their benefits and drawbacks, and how they are used in various food matrices. We will also go over the difficulties in applying MS-based techniques to routine food safety monitoring as well as the potential of this technology to protect public health and global food security.

2. CONVENTIONAL METHODS FOR THE DETECTION OF FOOD TOXINS

There are several specific and well-established conventional methods used on a regular basis for food safety assessment and regulation. These methods are often specific, but they may have limitations in terms of sensitivity, speed, and ability to detect a wide range of toxins (Figure 1). Some of the most common conventional methods for the detection of food toxins are mentioned below.

2.1. Immunoassays. The most commonly used immunoassay methods in the field of food toxin detection are enzyme-linked immunosorbent assays (ELISA) and lateral flow immunoassays. These methods are very rapid and help with easy detection. Samples are allowed to interact with either labeled enzymes or antibodies, thus helping in the detection. However, compounds with similar core structures or nontoxic analogs can provide false positive results. Phycotoxins like Okadaic acid, Yessotoxin, Pectenotoxin, Azaspiracid, Cyclic imines, Palytoxin, Domoic acid, Saxitoxin, Microcystin, and Cylindrospermopsin; mycotoxins like Aflatoxin B1, Deoxynivalenol, Fumonisin B1 Zearalenone, and T-2; and bacterial

toxins like *Clostridium perfringens* α , β , and ϵ toxin, *Staphylococcal* enterotoxins A, B, C, and E, botulinum toxins, and *Escherichia coli* enterotoxins^{49–51} are detected using immunoassay methods.

2.2. Chromatographic Techniques. Thin-layer chromatography (TLC), high-performance liquid chromatography (HPLC), and gas chromatography (GC) are commonly used to separate and quantify food toxins. Among various seafood-originated toxins, Domoic acid, paralytic shellfish toxins, and Aflatoxin B1 can be easily detected by HPLC and TLC.^{52–55} Among all, TLC is a simple technique in which food toxins are chromatographed on a plate layered with a thin layer of stationary phase using different mobile phase solvents, aiding the separation and visualization of the toxins. On the other hand, HPLC and GC are more advanced ways to separate toxins depending on various principles, enabling us to separate and quantify different food toxins. HPLC and GC analysis needs often include a complex protocol for sample preparation, which is one of the major limitations in the application of these techniques.

2.3. Spectroscopic-Based Techniques: UV–visible and Fluorescence Spectroscopy. Each toxin possesses a different core structure with different motifs and thus absorbs light at a particular wavelength, enabling its detection if monitored at their respective wavelength. This principle is harnessed for the detection of food toxins such as aflatoxins and can be measured with UV-fluorescence spectroscopy.⁵⁶ It was suggested that if a sample is showing response at 400 and 550 nm with respect to 365 and 730 nm excitation wavelengths, it is supposed to be contaminated with aflatoxins. Singh et al.⁵⁷ also reported that aflatoxin B1 and ochratoxin A have maximum absorption (λ_{max}) at 365 and 380 nm. Both UV–visible and fluorescence spectroscopic techniques are easy to handle and cost-effective. Moreover, fluorescence spectroscopy has a high sensitivity for the detection of food toxins. Despite these advantages, there are certain toxins whose absorption and emission wavelengths may not be very selective and specific, making one of the major limitations in their detection by this technique. Other spectroscopic methods like nuclear magnetic resonance (NMR) spectroscopy are also used to elucidate the complex structure of food toxins.⁵⁸ Similarly, near infrared (NIR) spectroscopy uses NIR (14000–4000 cm^{-1}) wavelength that causes vibration of C–H, O–H, N–H, and C=O bonds in the biomolecules and can be helpful for the detection of toxins in the food. Recently, this technique was used to detect Diarrhetic shellfish toxins in the mussels.⁵⁹

2.4. Biological Assays. One of the most conventional methods is the direct injection of toxic samples into live animals and monitoring of their physiological response, behavior, and mortality. This assay is commonly used for the detection of marine toxins like Diarrhetic shellfish toxins in seafood.^{60,61} However, these tests are time-consuming and costly and raise major ethical concerns.

2.5. Biochemical Assays. Biochemical assays do not detect food toxins by direct measurement but rather involve the measurement of toxin-induced biochemical changes such as enzymes. For example, the inhibition of phosphatase activity is used to detect the presence of diarrhetic shellfish toxins.^{62,63} The use of this technique is very narrow, utilizing different instruments and reagents, and cannot be applied to multiple types of toxins, thus making it less applicable for the detection of various food toxins.

2.6. Microbiological Assays. Some food toxins are produced by certain microbes. Hence, assays involving the presence of the microbe are sometimes used as a proxy to detect their presence. For instance, the detection of *Bacillus cereus* in the food samples can point toward the presence of enterotoxins.^{64,65} Although these assays are easy to operate and inexpensive, they lack sensitivity and specificity.

2.7. Sensor-Based Approaches. Sensor-based methods of toxin detection in foods are very popular since they can be used on the site. There are many toxins such as aflatoxin B1, diarrhetic shellfish toxins, and microcystins that can be detected by sensors.⁶⁶ Examples may be biosensors and aptasensors. Undoubtedly, the use of sensors helps to screen samples for the possible presence of toxins. However, the sensors may vary in terms of sensitivity and specificity.

3. MASS SPECTROMETRY-BASED DETECTION

Mass spectrometry (MS) is a powerful analytical technique used to detect and quantify various compounds including food toxins. This method can provide highly sensitive and specific results, making it a valuable tool for food safety and quality control.⁶⁷ Infectious toxins like prions and Shiga toxins can also be analyzed using mass spectrometry, where peptides are digested by proteases and then the digested proteinaceous parts are analyzed by MS.⁶⁸ The pervasive contamination of food products with mycotoxins has made monitoring their levels essential. Detection of mycotoxin biomarkers in urine provides valuable and specific data for exposure assessment to these food contaminants in order to overcome the disadvantages of the indirect approach based on food analysis.⁶⁹ Due to the diverse chemistry and occurrence of food toxins in feedstuffs and foods with complex matrices, the detection has become difficult. The primary source of error in the analysis results from inadequate sampling and inefficient extraction and cleaning procedures. Gas chromatography (GC)-MS is used to analyze volatile and semivolatile compounds, such as certain mycotoxins and pesticide residues, in a variety of dietary products. Before entering the mass spectrometer for ionization and detection, the compounds are vaporized and separated according to their volatility.⁷⁰ The principle of detection of food toxins using GC-MS includes multiple target analyte extraction using multi-residue analytical methods like QuEChERS and adsorption extraction.⁷¹

Recently, mass spectrometry has become one of the most effective methods even for identifying specific microorganisms by using matrix-assisted laser-desorption time-of-flight (MALDI-TOF) MS, followed by recognition of MS spectra unique to that organism that create a reliable fingerprint.⁷² MALDI-TOF MS-based identification of bacteria is more rapid, accurate, and cost-efficient than conventional phenotypic techniques and molecular methods. Rapid and reliable identification of food-associated bacteria is of crucial significance for product quality. In contrast to genotyping methods, it can also be readily implemented in routine analysis. Due to short turnaround periods, low sample volume requirements, and low reagent costs, MALDI-TOF MS has recently emerged as a powerful tool for the identification of food toxins or toxin-producing microorganisms.⁷³ Food toxins from various fields of seafood, fruits, vegetables, milk, dairy products, and oils can be detected using MALDI-TOF MS.⁷⁴ The microbial databases with unique features relevant to each microbial species are the key components and are therefore continually building up in size

Table 2. Variants of Mass Spectrometry (MS) and Their Advantages and Disadvantages

Technique(s)	Important features	Advantages	Disadvantages	Ref
Accelerator mass spectrometry (AMS)	<ul style="list-style-type: none"> Employed particle accelerator technology into a mass spectrometer. Its detection range include ion currents of more abundant stable isotopes (e.g, 12C, 13C) to very rare radionuclides (e.g, 14C) Simple and robust technique for regular analysis Able to detect nonvolatile compounds like sugar and proteins that cannot be detected in GC-MS 	<ul style="list-style-type: none"> Small quantity of sample is sufficient Need less time for estimation 	<ul style="list-style-type: none"> High cost makes it less affordable Small sample size makes it prone to contamination 	82, 83
Liquid chromatography mass spectrometry	<ul style="list-style-type: none"> Sample exposed to high temperature Used for the detection of volatile compounds from sample Nonvolatile compounds like sugar can be detected after derivatization 	<ul style="list-style-type: none"> Wide linear dynamic range Lower detection limit High precision and accuracy Analysis is faster and selectivity, Lower detection limits 	<ul style="list-style-type: none"> Lower accuracy Isotopes cannot be detected 	84, 85
Gas chromatography mass spectrometry	<ul style="list-style-type: none"> Good for the identification of unknown samples Efficient for nontargeted analyses 	<ul style="list-style-type: none"> Analysis is faster and selectivity, Lower detection limits 	<ul style="list-style-type: none"> Destructive method of analysis Only thermolabile compounds can be detected 	86–88
High resolution mass spectrometry	<ul style="list-style-type: none"> Traditional method for the identification of microorganisms Sample is first ionized, and segregated based on mass-to-charge ratio Measurement is done by determining with time-of-flight Used to measure the element level in sample 	<ul style="list-style-type: none"> Highly accurate and selective measurement Able to detect mass accurately with even small change is detectable Reduced analysis time 	<ul style="list-style-type: none"> Expensive analysis Data generated is huge and complex Not suitable for regular analysis of known samples In some cases, unable to differentiate between closely related species, e.g., <i>E. coli</i>, and <i>Shigella</i> Lack sufficient spectra in database 	89
Matrix-assisted laser desorption-ionization time-of-flight mass spectrometry	<ul style="list-style-type: none"> Sample converted to aerosol from liquid 	<ul style="list-style-type: none"> Wide analytical range with lower detection limit Need small quantity of sample High throughput with multielement detection 	<ul style="list-style-type: none"> High cost of investment and operation Need experts for operation 	90
Inductively coupled plasma mass spectrometry	<ul style="list-style-type: none"> It is also known as SELDI-TOF-MS-based ProteinChip System It is modified form of MALDI-TOF Proteomic profiling of biological fluids Two or more MS units are interconnected with quadrupoles and TOF analyzer Effective in analyzing complex mixture 	<ul style="list-style-type: none"> High throughput with multielement detection Less time-consuming and high through put system Highly specific 	<ul style="list-style-type: none"> Low detection precision for individual proteins from complex Low mass resolution High operational cost Limited sample through put 	91
Surface-enhanced laser desorption/ionization time-of-flight mass spectrometry		<ul style="list-style-type: none"> Low signal-to-noise ratio Able to detect covalent modifications in proteins Sensitive and reproducible 		92, 93
Tandem mass spectrometry				79, 94, 95

with the updated information on newly discovered microbial species and their annotations.

Another powerful MS-based tool routinely used for food toxin detection is liquid chromatography (LC)-MS due to its advantages in terms of sensitivity and selectivity. LC-MS is widely used for the analysis of mycotoxins, alkaloids, marine toxins, glycoalkaloids, cyanogenic glycosides, and furocoumarins in food. The excellent sensitivity, even at low concentration levels, selectivity, and capacity to resolve coeluting compounds based on their molecular masses make LC-MS currently the most effective technique for the simultaneous detection of multiple regulated, unregulated, and emerging toxins in a single run. Commonly, the LC-MS methods for the quantitative determination of natural toxins are based on the use of a triple-quadrupole analyzer, tandem mass spectrometry, and multiple reaction monitoring (MRM) modes. The co-occurrence of natural toxins in combination with other chemical contaminants, such as pesticides, growth regulators, and veterinary drugs, as well as bioactive compounds (i.e., lignans, flavonoids, and phenolic compounds), in a wide variety of food matrices has increased the demand for analytical methods addressing the simultaneous determination of multiple analyte classes. LC-MS method used for the detection of food toxins includes alkaloids, furocoumarins, cyanogenic glycosides, marine toxins, and mycotoxins.⁴¹

Inductively coupled plasma mass spectrometry (ICP-MS) is a powerful analytical technique for the detection of elements like heavy metals that allow multielement detection simultaneously with high speed and at very low concentrations.⁷⁵ Further, for the enhanced selectivity, inductively coupled plasma mass spectrometry (ICP-MS) and tandem mass spectrometry (MS/MS) can be utilized. In the ICP-MS technique, the sample under examination is digested by employing a suitable technique such as dry ashing, acid digestion, or microwave digestion, etc. to solubilize the analytes of interest. Further, the sample is injected into an inductively coupled plasma source which ionizes the sample and detected by MS.⁷⁶ Ion mobility spectrometry (IMS) is another advancement in analytical techniques that relies on ion mobility that is under the influence of velocity of ions and strength electric field.^{77,78} In order to take advantage, multiple MS-based approaches can be merged and operated together, which improves the efficiency of analytical techniques. Tandem mass spectrometry (TANDEM MS), also called MS/MS, is one such approach in which samples are analyzed either by multiple mass spectrometers connected to each other or with different analyzers arranged sequentially.⁷⁹ A MS technique can also be coupled with immunoaffinity chromatography, and it is known as IAC-MS. This technique uses antibodies-based columns to acquire selectivity and also to isolate target analytes from the sample matrix.⁸⁰ The isolated target molecules are injected into the MS component, which provides high sensitivity and identification of toxins based on mass-to-charge ratio. IAC-MS provides exceptional selectivity and can be used for the detection of various types of food toxins such as mycotoxins, pesticides, veterinary residues, and allergens, etc.⁸¹ MS-based approaches have offered diverse and advanced modules for the detection of analytes precisely. Table 2 summarizes the advantages and disadvantages of MS-based approaches under different conditions.

4. RECENT ADVANCEMENTS IN FOOD TOXIN DETECTION USING MS

Food toxins have become a serious concern for the society. The presence of various types of contamination and toxins like pesticides, herbicides, microbial metabolites, and plant-based toxins is highly detrimental to human health even at ppb concentrations when present in food, water, or animal feed.⁹⁶ Safe food is explicitly a matter of concern and is indispensable for human health. In the current scenario, an effective and sensitive detection method becomes necessary to detect contamination of food and water with chemicals and pathogenic microbes and related products. Existing conventional methods for food analysis, based on PCR, chromatography, and spectrophotometry, have shown significant reliability and accuracy; however, the cost of analysis, time consumption, and requirement of specialized personnel impede their usage for frequent monitoring of food samples (as already summarized above). Hence, there exists an upsurging thrust for innovating rapid, accurate, robust, but inexpensive alternatives for in situ and real-time detection of contamination of food samples. The primary step involved in every methodology to identify the food toxins in the sample under examination involves the extraction of food toxins from the matrix followed by purification to remove other substances that can interfere with the analysis.⁹⁷ After successful extraction of toxins, the sample is ionized into ions with ionization techniques such as electrospray ionization (ESI), chemical ionization (CI), or APCI (atmospheric pressure chemical ionization) or desorption techniques such as matrix-assisted laser desorption ionization (MALDI). These ions from the ionization chamber are accelerated followed by deflection in the magnetic field due to a difference in their masses. The beam of ions is then analyzed by the detector based on their mass-to-charge ratio. There are many types of mass analyzers available such as magnetic sector analyzers, quadrupole mass analyzers, double focusing analyzers, time-of-flight analyzers, etc. Further, MS can also distinguish between different food toxins depending upon their mass-to-charge ratio. In order to overcome the limitations associated with the conventional methods, MS can be coupled with these techniques to enhance the capabilities in complete and accurate analysis of different food toxins present in the sample.⁹⁸ GC technique relies on the comparison of retention times with known standards and also lacks in the distinction of structurally similar compounds. Thus, to enhance selectivity, identification, and elucidation of the structure of various toxins present in the sample under examination, GC is coupled with the MS.⁹⁹ HPLC technique requires optimization such as the selection of columns and mobile phases for each specific class of toxins. In addition to this, the sensitivity of HPLC is also low as that of MS techniques. HPLC techniques are more time-consuming than MS as they involve complex procedures for sample preparation.¹⁰⁰ We are herein summarizing various types of food toxins and the recent advancements in their detection using MS.

4.1. Mycotoxins. Mycotoxins are toxic secondary metabolites produced by fungi. These toxins can accumulate in the fungal-contaminated food grains such as corn, cereals, and legumes, and upon ingestion can traverse into the food chain affecting humans and animals.¹⁰¹ As per the Rapid Alert System for Food and Feed of EU (RASFF) report, mycotoxins contaminate around one-quarter of global food grain production both during pre- and postharvest.¹⁰² This clearly indicates the severity of the mycotoxin problem in the food that we consume.




Figure 2. General structures of some of the Aflatoxin variants.




Figure 3. General structures of various mycotoxins.

In the literature, around 300 mycotoxins are reported, but only seven toxins are quite common in food worldwide such as aflatoxins (AF), trichothecenes (TC), zearalenone (ZEN), fumonisins (FB), ochratoxins (OTA), citrinin (CIT), and

patulin (PAT).¹⁰¹ Fungal species belonging to the genera of *Aspergillus*, *Penicillium*, and *Fusarium* are predominantly toxigenic and most frequently lead to cases of mycotoxin concentration.¹⁰¹ AF (B1, B2, G1, and G2) produced by

Table 3. Regulatory Permission Limits for Various Mycotoxins in Food and Feed Products As Per European Commission^{122–127}

Category	Food or animal feed products	Permissible limit ($\mu\text{g}/\text{kg}$)	Category	Food or animal feed products	Permissible limit ($\mu\text{g}/\text{kg}$)
Aflatoxins B1 (AFB1)			Aflatoxins (AFs) total		
Food	Brazil nuts, groundnuts, hazelnuts, and oilseeds for human consumption after physical treatment	8		Maize and rice (direct or as ingredients) for human consumption after physical treatment	10
	Almonds, apricot kernels, and pistachios for human consumption after physical treatment	12		Citrinin (CIT)	
	Brazil nuts, groundnuts, hazelnuts, and oilseeds for human consumption directly (No physical treatment)	2–5	Food	Food supplements prepared from red yeast fermented rice	2000
	Almonds, apricot kernels, and pistachios for human consumption directly (No physical treatment)	8		Deoxynivalenol (DON)	
	Dairy products for consumption by infants, baby food, and processed cereal-based food	0.1	Food	Unprocessed durum, maize, oats and wheat	1750
	Spices	5		Cereals and cereal flour for direct human consumption	750
	Dried fruits, and Figures for human consumption after physical treatment	5–6		Cereal-based processed foods and baby food	200
	Dried fruits (except Figures) for human consumption directly (No physical treatment)	2	Feed	Animal feed from—cereals	8–10
	Maize and rice (as ingredients) for human consumption after physical treatment	5		Complete as well as complementary feeding stuff	0.9–5
Feed	Feed materials	0.02–0.05		Fumonisin (FB1+FB2)	
	Complete feeding stuff with the exception of Calves, cattle, and lambs	0.05	Food	Unprocessed maize	4000
	Poultry	0.005–0.01		Maize for direct human consumption	1000
	Complementary feeding stuff	0.005–0.05		Maize-based processed food for babies and young children	200
Aflatoxins M1 (AFM1)			Feed	Maze based feed	60
Food	Milk	0.05		Complete and complementary feeding stuff	5–50
	Infants' dairy products, baby formula and baby milk,	0.025		Ochratoxins (OTA)	
Aflatoxins (AFs) total			Food	Cereals-based unprocessed products	3–5
Food	Almonds, apricot kernels, brazil nuts, groundnuts, hazelnuts, oilseeds, and pistachios for human consumption after physical treatment	15		Cereal-based processed food and baby food, dietary foods products specially for medical purposes	0.5
	Groundnuts, oilseeds, and processed products for human consumption directly (No physical treatment) or as ingredient	4		Beverages based on grapes	2
	Almonds, apricot kernels, brazil nuts, hazelnuts, pistachios, and for human consumption directly (No physical treatment)	10		Coffee roasted/instant	5/10
	Spices	10		Spices	15
	Dried fruits, and Figures for human consumption after physical treatment	10	Feed	Cereals-based feed materials	0.25
	Dried fruits (except Figures) for human consumption directly (No physical treatment)	4		Complete and complementary feeding stuff for poultry	0.1
				Patulin (PAT)	
			Food	Fruit juices	50
				Solid apple products	25
				Solid apple as well as apple juice for babies and young children	10
				Zearalenone (ZEN)	
			Food	Cereal products (unprocessed except maize)	100 (350)
				Cereals for direct human consumption	75
				Maize for direct human consumption	100
				Cereals and maize processed products for babies and young children	20
			Feed	Cereals and maze-based feed materials	2–3

Aspergillus), CIT (by *Penicillium*, *Aspergillus*, and *Monascus* etc.), ergot alkaloids (*Claviceps purpurea*), FB (*Fusarium* sp.), ochratoxin A (produced by *Penicillium* and *Aspergillus*), PAT (*Penicillium patulum*), TC (*Fusarium*, *Stachybotrys*, *Trichothecium*, and *Trichoderma* sp.), and ZEN (*Fusarium graminearum*) are some of the well-known mycotoxins responsible for serious lethal reactions like cancer induction, kidney toxicity, immune suppression, stachybotryotoxicosis, turkey X syndrome, etc.¹⁰³ Another issue with these mycotoxins is their resistance and tolerance toward thermal treatment; hence, they remain active even after heating.¹⁰⁴

AFs are groups of potentially toxic fungal secondary metabolites reported from *Aspergillus flavus*, *A. parasiticus*, and *A. nomius* (Figure 2). They are produced from polyketides and commonly present in cereal crops like corn, peanuts, walnuts, wheat, etc.¹⁰⁵ AFs can lead to chronic toxicity related to hepatic tissues, necrosis, hepatomas, periportal fibrosis, jaundice, hemorrhage, and fatty liver changes, and also exhibit teratogenicity, carcinogenicity, and immunotoxicity.^{105,106}

They were first reported in the state of Gujrat and Rajasthan, India in 1974, which resulted in the onset of hepatitis caused by *A. flavus* infected staple food and maize. *A. flavus*, *A. parasiticus*, *A. nomius*, and sometimes *Emericella* spp. are the main producers of AFs.¹⁰⁵ To date, more than 20 AFs variants have been reported, among which variant B1 is the most common and most lethal. AFs B1 and B2 are produced by *A. flavus* and *A. parasiticus*, and AF M1 is produced by *A. parasiticus* and can be transmitted through milk. Variants M1 and M2 are also produced by metabolism of B and B2.¹⁶ AF B1 is metabolized by the P450 monooxygenase system and generates AF 8,9-epoxide (reactive epoxide) that induces mutations and cancer by forming DNA adducts.¹⁰⁶

Besides *Aspergillus*, isolates belonging to *Fusarium* are also known to produce the most potent toxins, which include deoxynivalenol (DON), FBs, and ZENs (Figure 3). DON belongs to the sesquiterpenoid group of trichothecenes. It mainly contaminates corn, wheat, and barley. It is also a toxic secondary metabolite that has a negative health impact on the

Table 4. Detection of Mycotoxins by MS-Based Approach

Name	Sample	Method	Operating parameters	Outcome	Ref
Trichothecenes T2	Wheat and maize	Portable mass spectrophotometer	MS parameter: heater 200 °C; sample pump 0%; mass center: $m/z = 484$; mass tolerance: 1; spectrum average: 10	Limit of detection 0.2 mg/kg	131
Alternaria toxin	Fruits and vegetables	QuEChERS extraction followed by MS analysis	Triple quadrupole MS; pressure curtain gas 35 psi and collision gas 8 psi; ion spray voltage 5500 V (positive ion mode), and -4500 V (negative ion mode)	Detection limit 1.0–5.0 $\mu\text{g}/\text{kg}$ (Extraction recoveries 73.0–120%) Repeatability <12.9%	132
Mycotoxins	Standard sample	UHPLC-QTrap-MS/MS	Injection volume: 20 μL ; column temperature: 30 °C; flow rate: 0.2 mL min^{-1} ; mobile phase 0.1% formic acid+ acetonitrile run time: 25 min	Limit of quantification 0.005 and 13.54 ng mL^{-1} (Limit of detection 0.001–9.88 ng mL^{-1} ; 0.005; Recovery 67.5–119.8%)	133
Aflatoxin B1 and M1	Blood	HPLC–MS/MS	Allure PEP column; positive electrospray ionization; source temp 500 °C	LOD 0.05 to 0.2 ng/mL; accuracy 92–111%	134
Alternaria toxins	Vegetable sample	UPLC–MS/MS	ACQUITY UPLC BEH C18 column; injection volume: 5 μL ; column temperature: 40 °C; flow rate 0.4 mL min^{-1} ; mobile phase: 0.1% formic acid water; gradient flow 10–90%	Contaminated solanaceous vegetable: 41.1% AME; 4.26%; AOH: 6.38%; altenuene: 6.38%; tentoxin: 42.6%; tenuazonic acid: 55.3%	135
Aflatoxin M1		HPLC–MS/MS	Quantitative daughter m/z : 273; qualitative daughter m/z : 259; collision energy 23 eV	Quantification limit 1.62 ppb; detection limit 0.54 ppb; 98.5% accuracy	136
Ochratoxin A	Coffee and tea	UHPLC–MS/MS	Triple quadrupole MS; electrospray ionization, ion source temp 300 °C; flow rate 3 L/min	Sensitivity 0.30 and 0.29 ng/mL	137
Trichothecenes	Oat based products	U-HPLC–HRMS/MS	Reverse phase column; electro-ionization detector; column temperature 40 °C; ethanol +water+formic acid gradient elution	Frequency of free T2 toxin 92% T2 mono glucoside 69%	138
Alternaria toxins	Rice	LC–MS/MS	Separation with hyper clone column and detection with C18 column; detection with negative electrospray ionization, source temperature 300 °C	Limit of detection and quantification AME: 0.03 and 0.09 $\mu\text{g}/\text{L}$; altenuene: 5.48 and 16.24 $\mu\text{g}/\text{L}$	139
Alternariol monomethyl ether (AME), Alternariol (AOH), and tentoxin	Standard sample	LC-ESI-MS/MS	LC–MS/MS with triple quadrupole; separation at 25 °C with C18 column; analysis with quadrupole MS, source temperature 450 °C; ion spray voltage 5500 V	Limits of detection and quantitation: 0.7 and 3.5 ng/g recovery 80%	140

consumer by disturbing the intestinal barrier and has immunostimulatory as well as immune-suppression properties at low and high doses, respectively.¹⁰⁷ The same group also contains its acetylated derivatives named nivalenol, T-2 toxin, and HT-2 toxins.¹⁰⁸ After ingestion, DON is absorbed and metabolized in the intestine via DON-3S, DON-GlcA, and DOM-1. In poultry birds, DON-3S and DON-15S are eliminated via bile and urine, while in swine, it is absorbed in the upper digestive system; hence, poultry birds are the least sensitive to these toxins, and swine are most sensitive to these toxins. In humans, contaminated foods like infected meat and cereals are the most common sources.¹⁰⁷ *Fusarium* sp. is also responsible for other mycotoxins named FBs (secreted by *Fusarium verticillioides* and *Fusarium proliferatum*). *Aspergillus niger* is also able to produce FBs. Such toxins are commonly reported from cereals like peanuts, maize, rye, oats, millets, and grape.¹⁰⁹ To date, more than 15 homologous forms of fumonisin have been reported that are referred to as A, B1, B2, B3, C, P, etc. However, B1 is the most toxic form of FB.^{110,111} FBs are known to have carcinogenic, neurotoxic, and hepatotoxic effects that cause hepatocarcinoma, defects in the neural-tube, and nephrotoxicity.¹⁰⁹ ZEN is a nonsteroidal, estrogenic mycotoxin produced by *F. acuminatum*, *F. crookwellense*, *F. culmorum*, *F. cerealis*, *F. equiseti*, *F. graminearum*, *F. oxysporum*, *F. sporotrichioides*, *F. semitectum*, and *F. verticillioides*.¹¹² It disrupts reproductive capacity by affecting mammalian folliculogenesis and impairs granulosa cell development and follicle steroidogenesis.¹¹³ It is thermostable¹¹⁴ and resistant to processing stress like milling and storage.^{115,116} ZEN leads to kidney damage and liver injury and causes inflammation.¹¹²

Ergot alkaloids are toxic secondary metabolites produced by several fungi of *Clavicipitaceae* (*Epichloë*, *Claviceps*, *Balansia*, and *Periglandula*) and *Aspergillus fumigatus*. The producers mainly include *Epichloë endophytes*, *Epichloë festucae* var. *lolii*, *Epichloë coenophialum*, and *Claviceps purpurea*.¹¹⁷ They belong to compounds containing an indole group and are derived from L-tryptophan. They cause “ergotism” and their toxic effect is reflected by hyperglycemia, gastrointestinal upset, mydriasis,¹¹⁸ and even endocrine disruption.¹¹⁹ Besides these major mycotoxins, some other chemicals including OTA and PAT have also been reported from fungi. OTA is nephrotoxic and produced by a diverse range of fungi such as *Aspergillus ochraceus*, *A. carbonarius*, *A. niger*, and *Penicillium verrucosum*. It can also lead to renal tumors, Balkan endemic nephropathy, and chronic interstitial nephropathy.¹²⁰ Patulin is produced by *Penicillium*, *Aspergillus*, and *Byssoschlamys*, while alternariol (AOH) and alternariol monomethyl ether (AME) are *Alternaria* toxins, produced by fungi of the *Alternaria* genus found in fruits and related products.²⁰

To ensure food safety with special consideration to mycotoxins, international, national, and regional agencies like the World Health Organization, Food Agriculture Organization, Codex Alimentarius Joint Expert Committee for Food Additives and Contaminants, European Food Safety Authority, GCC Standardization Organization, and Japanese Association of Mycotoxicology have determined the permission limit for mycotoxins contamination in food as well as feed.¹²¹ Table 3 summarizes the permissible limit for various mycotoxins in food^{122–125} and animal feed^{126,127} as per the European Commission.

HPLC and GC are the common approaches for the detection and higher accuracy and sensitivity making MS an elegant and dominant analytical tool for toxicological and metabolite

analysis.¹²⁸ In addition, it also allows simultaneous detection of a diverse range of toxins together and aids in method standardization and implication to ensure the rapid evaluation of samples for food safety analysis. Areo et al.¹²⁹ have employed UHPLC–MS/MS for the detection of AFs, ZEN, and OCT A from 100 tea samples (collected from registered shops within South Africa), prepared in acetonitrile/water/acetic acid solvent by QuEChERS extraction method. The supernatant was mixed with 900 mg of anhydrous MgSO₄, 150 mg of C18, and 150 mg of primary secondary amine that separates the organic phase. The organic phase was collected and dried under a nitrogen stream and further reconstituted in methanol/water for analysis via UHPLC–MS/MS. The method selected has very high linearity (>0.99) and precision (6–29%). AFs, i.e., AFB1, AFB2, AFG2, OCT A, and ZEN, were absent in the samples, and AFG1 was present in very low amounts 1.72–5.19 µg/kg that were also below the regulated level in food as recommended by EU Commission Regulation 1881/2006.¹²⁹ You et al.¹³⁰ have evaluated the effect of culture medium for mycotoxin accumulation by *Alternaria*. Secondary metabolites were characterized by nontarget analysis with HRMS. Mycotoxins produced by *Alternaria* were grouped into for families: alternariol monomethyl ether (AME), alternariol (AOH), altenuene (ALU), Desmethyl dehydro altenusin (DMDA), and dehydroaltenusin (DHA) families, Altertoxin-I (ATX-I) family, tentoxin (TEN) family, and tenuazonic acid (TeA) family. Culture medium greatly influenced the type of mycotoxins produced. *wiz* Potato Sucrose Agar medium is suitable for AOH, AME, ALU, ALT, DHA, and DMDA, while Potato Dextrose Agar supported the accumulation of ATX-I, TEN, and TeA. Table 4 summarizes some of the research for the detection of various mycotoxins with MS-based tools.

4.2. Bacterial Toxins. Bacterial contamination has shown diverse causes as *Shigella* mainly infect via unwashed hands, while *Campylobacter* and *Escherichia coli* are usually present in raw milk, undercooked meat and poultry products, and contaminated water. In contrast, *Listeria monocytogenes* and *Yersinia enterocolitica* are found in refrigerated food.¹⁴¹ Bacterial toxins have been classified as endotoxins and exotoxins. Structurally, endotoxins have distinct structural regions, i.e., glycolipid is made up of disaccharide and fatty acids which are usually capric, lauric, myristic, palmitic, and stearic acids. These acids are buried within the outer cell membrane of the bacterium. The nucleus is the second important part, which is made up of a hexose- and heptose-based heteropolysaccharide. The glycolipid and nucleus are interconnected by the sugar acid 2-keto-3-deoxyoctanate. Endotoxins are lipopolysaccharides and are part of the outer membrane of Gram-negative bacteria. These are also identified as important determinants and antigenic parts of bacteria that aid in attachment with the host as well as in pathogenicity. Exotoxins are proteins in nature that are released by Gram-negative bacteria and disrupt cell division, causing lysis and tissue damage.^{142,143} Exotoxins are further classified into types I, II, and III based on the mechanism of action. Toxin type I can make critical changes in the host's cells without internalizing. Superantigens secreted by *Staphylococcus aureus* and *Streptococcus pyogenes* are examples of a type I toxin. The type II group includes hemolysins, phospholipases, aerolysin, and GCAT proteins. It intrudes the host cells and creates pores to destroy the host cell's membrane. In comparison to type I and II, type III is a diverse group in terms of activity. It has a binary structure with fractions A and B. Fraction B in the toxin facilitates the binding with receptor in the host cell, while




Figure 4. General structures of various marine biotoxins.

another fraction, i.e., “A” carries enzymatic activity and is responsible for the toxin effect. Anthrax toxin (*Bacillus anthracis*), Cholera toxin (*Vibrio cholerae*), and Shiga toxin (*Escherichia coli* O157:H7) are some examples of exotoxins.^{142,144} Botulinum is 150 kDa and composed of a heavy chain of 100 kDa and a light chain of 50 kDa. Heavy chain is responsible for binding to receptors on neuron surface, while light chain cleaves proteins required for nerve signal transmission, i.e., botulinum A, C, and E cleave synaptosomal-associated protein 25, while B, D, F, and G variants act on synaptobrevin-2.^{72,145,146} In a similar fashion, *B. anthracis* produces three types of proteins or factors, e.g., lethal factor, edema factor, and protective antigen. Protective antigens split and form a fragment of 63 kDa that forms heptamers and octamers to finally find the cell surface. In addition, they also bind with lethal factors to form lethal toxin.^{72,147}

4.2.1. Detection of Bacterial Food Toxins. *Clostridium*, *Salmonella*, *Staphylococcus*, and *Listeria* are some common pathogens causing foodborne infections in humans. Previously used methods, e.g., enzyme immunoassay (ELA), were fast and sensitive, but their accuracy was limited due to cross-reactivity reporting a high rate of false positives and misguided public health care personnel. Techniques based on MS are powerful and can be multiplexed for the detection of various protein toxins, e.g., toxins from *Clostridium*,¹⁴⁸ *Bacillus*,⁸⁸ and many more with speed, sensitivity, and accuracy. Botulinum, a neurotoxin, is produced by *Clostridium botulinum* in seven different serotypes (A–G). Specific detection of these toxins from different strains needs high analytical sensitivity and a MS-based approach. The enzymatic activity-based approach relies upon substrate fragments generated by these toxins which can be used as targets by MALDI-TOF MS.⁷²

In 2002, a peptide mass map of toxin variants A1 and B1 was prepared by targeting the trypsin digest of the toxin by van Baar and colleagues. The work was extended to C, D, E, and F in

2004.⁷² In successive generations, several advancements have shown effective approaches like endopep-MS.¹⁴⁹ Rosen et al. have developed the endopep-MS-based method for the identification of botulinum A and E simultaneously and rapidly.¹⁵⁰ Both A and E identify the same target SNAP 25 protein but act on different sites. 3D structures of both types of fragments were used for differential identification. Drigo et al.¹⁵¹ employed the same endoPep-MS approach for botulinum toxins C and D. The method has shown a sensitivity of 100% with specificity and accuracy of 96.08% and 97.47%, respectively. Integration of MS with other analytical methods like HPLC, GC, FPLC, etc. has improved bacterial toxin profiling. Toxoflavin and fervenulin are bacterial toxins produced by *Berkholderia* and *Streptomyces hiroshimensis*. These compounds are common contaminants in fermented corn flour, rice bran oil, distiller’s yeast, sweet potato starch, *Tremella fuciformis* Berk., and rice noodles. These compounds are sensitive to degradation in 1% ammonia solution. UHPLC-Q-TOF/MS allowed for the detection of degradation products. The modified approach led to lower down the limits of detection of toxoflavin and fervenulin to 12 $\mu\text{g}/\text{kg}$ and 24 $\mu\text{g}/\text{kg}$, respectively, with recovery of 70.1–108.7%.¹⁵² The MS approach also employed natural phenomena of antigen–antibody interactions for the detection of toxins and related antigens. *Salmonella typhi*, Gram-negative enterobacteria, is responsible for typhoid fever and meningitis. The immunoreactive proteins of bacteria were used as targets to develop improved diagnostic tools with MS. An immunaffinity-based proteomic approach was employed with IgG and IgM antibodies from typhoid patients. The approach aided in the identification of 28 immunoreactive proteins, out of which 14 were complementary to IgG, 4 for IgM, and the rest 10 for both, hence retained by respective charged columns. In context to antigenicity, 22 proteins have shown antigenicity and immunogenicity.¹⁵³ Such an approach is helpful for rapid identification, and its reproducibility and reliability can be




Figure 5. General structures of various marine biotoxin (Brevetoxin).

employed for vaccine and drug development. Peptide mass fingerprinting technique (PMF) associated with MALDI TOF/MS or ESI/MS is a top-down MS protocol where proteins are directly ionized to create a fingerprint of individual proteins and applied for detection of various microbial strains. PMF of unknown organisms is compared to those existing in PMF databases or compared to the spectrum of biomarker proteins with the proteomic spectral database, using MALDI-TOF MS. Typically a mass range m/z of 2–20 kDa is used for species-level identification where ribosomal proteins representing 60–70% of microbial cells' dry weight and some housekeeping proteins are selected.¹⁵⁴ Thus, by comparing with extensive commercial databases, microbial contaminants can be traced to the genus, species, or strain level, and such an identification tool is conveniently adapted in diagnostic laboratories.¹⁵⁵ However, using biomarkers is not very common for identification since it requires prior insight into the genome sequences before creating the required databases for proteins' molecular masses. *Staphylococcus aureus* delta-toxin has been detected using whole-cell (WC) MALDI-TOF/MS and LC-MS to correlate the expression of delta-toxin with the status of the agr (accessory gene regulator) status. Mass spectra of pure toxin from wild type strains and mutants for agr-rnaIII gene were compared specifically at the position of the peak for delta-toxin.¹⁵⁶

Biosensors have taken up an important role in the accurate and fast detection of food contaminants even in very low concentrations¹⁵⁷ using biorecognition elements, such as antibodies, enzymes, nucleic acids, phages, etc., along with electrochemical, optic, or piezo-electric devices for the detection of food contaminants. MS-based biosensors are less prevalent as compared to optical or electrochemical ones^{158,159} but have the potential to overcome the drawbacks of conventional models of biosensors. A multitoxin biosensor-MS was developed for the detection of multiple bacterial toxins simultaneously. Biomolecular interaction analysis-MS (BIA-MS) that used a two-step

method, i.e., first bonding of toxin molecules to antibodies immobilized on a sensor chip using SPR (surface plasmon resonance) and then the bound toxin, was identified by MALDI-TOF/MS. The potential of the multiaffinity sensor chip was validated by the detection of endotoxin from *Staphylococcus* in mushroom and milk samples and it successfully detected multiple toxins at concentrations as low as 1 ng/mL.¹⁶⁰

4.3. Marine Biotoxins. Marine biotoxins are natural compounds released in the marine environment by algae and phytoplankton during harmful algal blooms (Figure 4). These compounds are highly toxic for consumers and not only are related to serious illness but also lead to the death of aquatic organisms and even humans.¹⁶¹ Due to continuous release in the surrounding environment, these biotoxins accumulate in aquatic and marine organisms such as mollusks and fishes. Based on the chemical nature and solubility, these biotoxins are hydrophilic and lipophilic. Hydrophilic biotoxins are water-soluble and can cause amnesic shellfish poisoning, paralytic shellfish poisoning, and emerging pufferfish poisoning, while other groups of lipid-soluble biotoxins are responsible for diarrhetic shellfish poisoning and azaspiracid shellfish poisoning. There is another group of toxins with less available information that is categorized as emerging toxins and can cause unregulated ciguatera fish poisoning, cyclic imines, and neurotoxic shellfish poisoning.^{162–165} Paralytic shellfish poisoning (PST) is a group of more than fifty-eight related compounds, produced by *Alexandrium* dinoflagellates of the Atlantic and Pacific coast and Mediterranean Sea. It has a tetrahydropurine skeleton among which saxitoxin (SXT) and gonyautoxin (GNT) are common. Structurally, STX has been categorized into four subgroups named carbamate, N-sulfo-carbamoyl, decarbamoyl, and hydroxylated saxitoxins.¹⁶⁶ The toxicity related to PST is reflected in mild as well as severe depending upon toxicity. The mild symptoms include numbness, tingling sensation around lips followed by expansion of the area, itching and prickly

Table 5. Detection of Bacterial and Marine Biotoxins by MS-Based Approach

Name	Sample	Method	Operating parameters	Outcome	Ref
Enterotoxin	Commercial	LC-HRMS	LC-MS Q ₂ -extractive mass spectrophotometer C18 reverse phase column; isocratic elution	93 signature peptides identified for enterotoxins	182
Botulinum	Commercial	Endopep MS	Triple quadrupole mass spectrometer; Turbo Ion Spray interface; C18 column; gradient elution	Toxin detection 0.1 MLD ₅₀ and quantification 0.62 MLD ₅₀	183
Okadaic acid	Raw and cooked food (Mussel, clam, flatfish)	Tandem mass spectrometry	C18 column with triple-quadrupole mass spectrometer; ammonium format gradient elution; negative ionization mode	LOD and accuracy 0.2–5.1 μg/k	184
Dinophysistoxin	Raw and cooked food (Mussel, clam, flatfish)	Tandem mass spectrometry	C18 column with triple-quadrupole mass spectrometer; ammonium format gradient elution; negative ionization mode	LOD and accuracy 0.2–5.1 μg/k	184
Anatoxins a (ATX) and Homoanatoxin-a (HAT)	Benthic-cyanobacterial-mat field samples	LC-HRMS/MS	Q ₂ Exactive HF Orbitrap MS; HESI-II electrospray ionization source; at 40 °C; resolution 60,000; collision energy 20 eV	Toxins are in conjugated form 15% ATX and 38% HAT	185
Cyanotoxins	Blue-green algae dietary supplement	Hydrophilic Interaction Liquid Chromatography-Tandem Mass Spectrometry	Electrospray ionization positive; source temperature 550 °C; ion spray voltage 5500 V; curtain gas 2.5 psi; collision gas 10 psi	Quantification limits 60–300 μg kg ⁻¹	186
Paralytic shellfish toxins	Marine shellfish	Hydrophilic interaction chromatography-tandem mass spectrometry	Separation with HILIC-Z column; acetonitrile and ammonium formate-formic acid as the mobile phase; positive electrospray ionization; samples are cleaned with by ion-pair SPE using a porous graphitic carbon cartridge	Limits of detection 1.7–13.7 μg kg ⁻¹ ; and quantitation 5.2–41.0 μg kg ⁻¹ ; recoveries 76.5–95.5%	187
Tetrodotoxin	Marine shellfish	Hydrophilic interaction chromatography-tandem mass spectrometry	Separation with HILIC-Z column; acetonitrile and ammonium formate-formic acid as the mobile phase; positive electrospray ionization; samples are cleaned with by ion-pair SPE using a porous graphitic carbon cartridge	Limits of detection 1.7–13.7 μg kg ⁻¹ ; and quantitation 5.2–41.0 μg kg ⁻¹ ; recoveries 76.5–95.5%	




Figure 6. General structures of various Phytotoxins. The general scaffold for Cyanogenic glycosides, Furocoumarin and Dehydropyrrolizidine, has been depicted with providing a few examples of various compounds belonging to the class of Furocoumarin.

sensation in fingertips and toes, dizziness, headache, and nausea. Moderate and severe illness symptoms include incoherent speech, prickly sensation and stiffness in limbs, weakness, difficulty in respiration, and muscular paralysis.¹⁶¹

Amnesic shellfish poisoning is mainly caused by domoic acid (DA) and derivatives produced by marine diatoms of *Pseudonitzschia*. DA, cyclic tricarboxylic amino acid, can bind with glutamate receptors in the central nervous system due to structural analogy and result in excess stimulation, induced production of reactive oxygen species (ROS), and ultimately cell death.¹⁶⁷ Consumption of DA resulted in gastrointestinal ailments including abdominal cramps, diarrhea, nausea, and vomiting. Neurological symptoms may also include confusion, disorientation, paresthesia, lethargy, short-term memory loss, and in severe toxicity cases, it may also result in coma or death.¹⁶⁸

Diarrhetic shellfish poisoning (DST) is a toxin produced by dinoflagellates of *Dinophysis* and *Prorocentrum*. It is a common type of contamination in shellfish industries due to over-extended prohibitions on mussel harvesting activity.¹⁶⁹ The responsible toxins for DST are a group of polyether compounds recognized as okadaic acid and its derivatives (dinophysistoxin); pectenotoxin; yessotoxin and its derivatives; and azaspiracid. Okadaic acid and azaspiracid consumption resulted in abdominal pain, diarrhea, nausea, and vomiting.^{161,170} Pectenotoxin and yessotoxin are not involved in human illness.¹⁷¹

Neurotoxic Shellfish Poisoning (NST) is another type of algal toxin that causes neurological as well as gastrointestinal ailments. Brevetoxins are a kind of marine biotoxin produced by *Karenia brevis* (Florida red tide dinoflagellate). It is a poly(ether ladder) compound (Figure 5). It causes mortality in massive fish and marine mammals. In humans, these toxins resulted in asthma-like symptoms if inhaled.¹⁷² Neurological and toxicity symptoms

of NST are paralysis, seizures, paresthesia, and coma, while gastrointestinal ailments are represented by nausea, diarrhea, vomiting, cramps, and bronchoconstriction, and extreme poisoning may also lead to death.¹⁶¹ Ciguatera fish poisoning (CFT) is one of the most common foodborne illnesses caused by marine biotoxin of ciguatera.^{173,174} Ciguaterins are toxic and lipid-soluble compounds found in marine organisms. Gambiertoxins, the precursor toxins, are produced by benthic dinoflagellates of *Gambierdiscus* genus. These toxins are accumulated in large predatory fishes like Spanish mackerels, moray eels, barracuda, and snappers.¹⁶¹ These compounds abnormally activate sodium ion channels and disrupt the cell membrane.¹⁷⁵ These compounds cause abdominal pain, nausea, diarrhea, vomiting, hypertension, and bradycardia along with neurological complications.¹⁶¹

4.3.1. MS Analysis and Detection of Algal and Marine Biotoxins. Algal and marine biotoxins are another class of heterogeneous toxins produced by algae and cyanobacteria, Dinoflagellates and diatoms produced during algal blooms in rivers, freshwater lakes, and marine aquatic systems.¹⁷⁶ Karunarathne et al., have found that 16,659 deaths have been reported in India between 1999 and 2018 due to poisoning.¹⁷⁷ In the case of sea food such as shellfish, exposure and contamination to multiple toxins are possible, hence an efficient system is able to detect diverse classes of toxins at the same time effectively. Blay et al.,¹⁷⁸ developed a method for the detection of multiple lipophilic biotoxins including azaspiracids, dinophysistoxins, and pectenotoxins as well as negative toxins via reversed-phase LC–MS within 7 min and hydrophilic toxins such as okadaic acid, dinophysistoxin-1,2, and yessotoxin from shell fish by recording scans at 2 Hz in positive and negative scans alternatively and 1 Hz in positive mode, respectively. Hydrophilic toxins including gonyautoxins domoic acid and saxitoxin

Table 6. Phytotoxins Detection from Food Samples via MS-Based Approach

Name	Sample	Method	Operating parameters	Outcome	Ref
Toxoflavin and Fervenuilin	Food samples and <i>Tremella fuciformis</i> Berk	UPLC-MS/MS	Separation column: ZORBAX SB-C18 column; oven temp: 35 °C; mobile phase: 0.1% formic acid+ methanol; flow rate 0.4 mL min ⁻¹	Detection limits (μg/kg) Toxoflavin 12 Fervenuilin 24 Toxin level (mg/kg) Toxoflavin: 7.5; fervenuilin: 3.2	152
Ricin	Soft drinks and serum	Surface-assisted laser desorption/ionization mass spectrometry	Pulsed Smartbeam II 2 kHz laser; wavelength 355 nm (~3.49 eV); frequency 1000 Hz; delayed extraction time 350 ns for proteins	Limit of detection 0.5 pmol/μL	195
Pyrrrolizidine Alkaloids	LC-MS/MS	Tea	UPHPLC with Quadrupole mass spectrometer; C18 Hypersil Gold column fitted; gradient elution	Total PA levels 13.4 to 286,682.2 μg/kg d.m	196
Ptaquiloside	LC-MS/MS	Bracken fern	LC-MS/MS; C18 column; gradient elution; column temperature 35 °C; electrospray positive ionization	Limits of detection 0.03 and quantification 0.09 μg/kg	197
Furanocoumarin	UPLC-MS/MS	Citrus sp.	Nexcol C18 column; column temp 40 °C; gradient elution; positive electrospray ionization	Compound recovery 94.07–114.53%	198
Amygdalin	LC-MS/MS	Kernels and Almonds	LC-MS/MS equipped with 6500 quadrupole linear ion trap (QTRAP) mass spectrometer and electrospray ionization	Recovery 90–107%; limit of detection 0. Ng/g and limit of quantitation 8 and 2.5 ng/g	199
Pyrrrolizidine alkaloids	quadrupole orbitrap MS	Honey	Polar C18 column; temp 40 °C; mobile phase flow rate of 400 μL min ⁻¹ MS detection with positive ionization mode; scan range 250–500 m/z and 70 k (fwhm)	Limit of quantification 0.1–0.3 μg kg ⁻¹	200
Pyrrrolizidine alkaloids	quadrupole orbitrap MS	Black and green tea	Polar C18 column; temp 40 °C; mobile phase flow rate of 400 μL min ⁻¹ MS detection with positive ionization mode; scan range 250–500 m/z and 70 k (fwhm)	Limit of quantification 1–11.7 μg kg ⁻¹	200
Pyrrrolizidine alkaloids	quadrupole orbitrap MS	Herbal infusion	Polar C18 column; temp 40 °C; mobile phase flow rate of 400 μL min ⁻¹ MS detection with positive ionization mode; scan range 250–500 m/z and 70 k (fwhm)	Limit of quantification 0.9–2.1 μg kg ⁻¹	200

were detected with mass accuracy of less than 1 ppm error and resolving power of 100,000 for the analytes (m/z 300–500). The limits of detection for lipophilic toxins were 0.041–0.10 $\mu\text{g/L}$ ppm (positive ions), 1.6–5.1 $\mu\text{g/L}$ (negative mode), and 3.4–14 $\mu\text{g/L}$ for domoic acid and paralytic shellfish toxins.¹⁷⁸ The biggest advantage of the method is that the analytes were detected with real time samples without any interference. Aquatic water bodies have a higher possibility of having aquatic biotoxins; hence, monitoring of water in water bodies is mandatory. Estevez et al. developed a method to seawater monitoring for marine biotoxins by hydrophilic interaction liquid chromatography coupled with HRMS. The main analytes considered for the detection were saxitoxin, decarbamoyl-saxitoxin, neosaxitoxin, gonaytoxin-2,3, and tetrodotoxin due to their adverse effects on gastrointestinal and central nervous systems in humans if taken up via seafood. Samples were processed via ultrasound-assisted solid–liquid extraction with methanol to extract toxins, followed by solid phase extraction using silica cartridges. The selected toxins are polar in nature; hence, the extraction stage is crucial for analysis, and the developed method has recoveries of 15–47% in filtrate and 26–71% in particulate fraction. Simultaneously, limit of detection was also affected with source as LOD was 0.5–5 $\mu\text{g/L}$ for filtrate and 3.1–62 $\mu\text{g/L}$ for particulate fraction.¹⁷⁹

Kolrep et al.¹⁸⁰ conducted comparative metabolite profiling to track the metabolism of okadaic acid in the liver and the role of CYP3A4 and CYP3A5 in its detoxification. It was found that LC–MS/MS can identify the metabolites distinctly from humans and rats based on the difference in +16 (+O) and +14 (+O/–H₂) Da. It suggested some critical differences in the metabolism of okadaic acid in humans and rats. In continuation, it was also found that rats generated more metabolites from okadaic acid in comparison to humans in the presence of NADPH-dependent enzymes.¹⁸¹ The establishment of metabolic patterns and fragments might be crucial for the identification of fingerprints for toxin identification. Table 5 elaborates on the detection of bacterial and marine biotoxins from different samples.

4.4. Phytotoxins. Phytotoxins are plant-derived compounds, including alkaloids and glycoalkaloids, that are naturally produced within plants but prove harmful if they remain in food products (Figure 6). These are secondary metabolites in plants and include cyanogenic glycosides, glucosinolates, glycoalkaloids, pyrrolizidine alkaloids, and lectins.¹⁸⁸ Based on the site, these toxins can be classified into endotoxins and exotoxins. Endotoxins may be normal metabolites that are present in cells but become harmful if consumed in higher concentrations, and these compounds are also referred as antinutritional factors, while exotoxins are toxic metabolites that are released from cells. Based on chemical nature, these are dehydropyrrolizidine, alkaloids, ptaquiloside, corynetoxins, and phomopsins.¹⁸⁹ Cyanogenic glycosides (CGLs) are present in almonds, cassava, bamboo roots, sorghum, and stone fruits. These toxins are generated from proteinogenic amino acids like leucine, isoleucine, phenylalanine, tyrosine, and valine as well as nonproteinogenic amino acids like cyclopentenylglycin. CGLs are potentially toxic for humans and result in acute cyanide intoxication, high respiration rate, lower blood pressure, headache, dizziness, stomach pains, diarrhea, vomiting, and mental confusion.^{188,189} Furocoumarins are found in many plants including carrots, celery roots, citrus fruits, parsley, and citrus plants. These compounds are responsible for gastrointestinal ailments and phototoxicity, skin reactions under UV

light.^{34,190} Lectins are reported from beans like kidney beans and can result in stomachache, diarrhea, and vomiting.¹⁸⁹

Phytotoxins are sometimes a part of the natural defense of plants, like Pyrrolizidine alkaloids (PAs) that are produced in *Asteraceae*, *Boraginaceae*, and *Fabaceae* families to defend plants against herbivores as well as insects. These toxins tend to have a common 1-hydroxymethyl pyrrolizidine core that is esterified with aliphatic acids. Besides edibles from these plants, honey is one of the common products contaminated with PAs, and hence, the extract or infusion can be used as an analyte for the detection of PAs. Ten plant samples, *Anchusa officinalis*, *Borago officinalis*, *Echium italicum*, *Eupatorium cannabinum*, *Heliotropium europaeum*, *Lithospermum officinale*, *Petasites hybridus*, *Senecio vulgaris*, *Symphytum officinale*, and *Tussilago farfara* from Orto botanicodella Scuola Medica Salernitana, Salerno, Italy, were collected, and aqueous extract was prepared with salting-out assisted liquid–liquid extraction. The aqueous extracts were analyzed with UPLC–MS/MS. The analysis was able to identify 88 PAs from 282 samples with an identification limit of 0.6–30 $\mu\text{g kg}^{-1}$ and a false negative rate <1.3% (at the concentration range of 4 $\mu\text{g L}^{-1}$).¹⁹¹

For simultaneous detection of phytotoxins and microbial toxins, HRMS was employed, which applied to over 156 compounds inclusive of about 90 plant toxins (e.g., various alkaloids and aristolochic acids), about fifty-four mycotoxins, and 12 phytoestrogens (e.g., lignan, isoflavones, coumestans, etc.) in plant-protein samples, like cereals. MS library, created on fragmentation pattern obtained with both negative and positive ionization modes for each toxin, using ten different collision energies was used for analysis. A typical workflow was followed with generic QuEChERS-like sample preparation, followed by UPLC using suitable mobile phases that allowed the resolution of over 50 toxic alkaloids. The method performance was evaluated for its sensitivity at levels ranging from 1–100 $\mu\text{g/kg}$, and reproducibility. The quantitation obtained against the standard addition approach could meet SANTE/12682/2019 criteria for 132 toxins out of the tested 156 toxin samples.¹⁹² The plant toxins ricin and RCA120 were detected, differentiated, and quantified by Kalb et al.,¹⁹³ via MS-based methods for the EQuATox proficiency test, in ~9 samples. They successfully identified the samples spiked with ricin or RCA120; samples spiked with a 0.414 ng/mL concentration could not be detected. Liang et al.¹⁹⁴ employed reversed phase LC–HRMS to detect five major phytotoxin groups including alkaloids, aromatic polyketides, flavonoids and steroids, and terpenoids at alkaline pH (>9). The developed method not only allowed the detection of 30 phytotoxins but also had forty-times higher detection sensitivity in comparison to older methods. Table 6 discusses some of the examples for phytotoxins detection using the MS-based approach.

4.5. Emerging Toxins. In addition to conventional toxins already known and summarized above, there is a group of toxic chemicals that are continuously evolving, mainly due to rising pollution. In the last eight years, the list of emerging chemicals has increased day by day. Synthetic chemical toxins include microplastics, organophosphorus and polybrominated flame retardants, perfluoroalkyl compounds, food process and packaging, waste substances, and nanomaterials.¹⁵ Besides, heavy metals, antibiotics and drug traces and metabolic intermediates, and agricultural chemicals²⁰¹ have shown bioaccumulation and have serious toxic effects if consumed, even in low concentrations. Health ailments include endocrine disruption, suppression and overexpression of the immune system,

inflammation, abnormal metabolic changes, skin diseases, carcinogenesis, etc., and the toxicity relies on interaction with the cellular system and receptors.^{15,201,202}

With the increase in pollution and intrusion of pollutants in the food web, the toxic chemicals traced in food and edibles are increasing. Some of those chemicals exhibited bioaccumulation and become silent killers, but some are potentially lethal. These emerging pollutants include pesticides, herbicides, healthcare, cosmetic chemicals, etc. Fipronil is a wide-spectrum phenylpyrazole insecticide used to control beetles, ants, cockroaches, etc. but its entry into the food chain is alarming due to its carcinogenic nature, necessitating its prohibition by the US Environmental Protection Agency (EPA). Suitable detection methods are thereby essential to identify and quantify these contaminants before they enter the food chain. The most reliable detection method includes LC–MS/MS and GC–MS, having specific sample preparation before the analyses (Table 7).²⁰³ One such preparatory method involved a modified QuEChERS sample preparation before using a triple quadrupole MS instrument coupled to ESI for detecting fipronil and its major metabolite fipronil sulfone, at concentrations of 5 µg/kg. The use of nontargeted approaches, such as SWATH-MS (sequential window acquisition of all theoretical mass spectra), enables the sequential analysis of fipronil and other such contaminants, e.g., pesticides and polyaromatic hydrocarbons. Glyphosate (insecticide) was detected in an underivatized form by innovating new extraction methods coupled with instrumentation. The QuPPE (Quick Pesticide Preparation) method was used for sample preparation²⁰⁴ followed by detection via sensitive MS instruments to achieve accurate quantitative results. LC–MS/MS was used in combination with the DMS (differential mobility separation) technique to terminate analytical interferences leading to improved signal by decreasing noise and, consequently, increasing accuracy and confidence in data. Using this method, LC–DMS–MS/MS was used for identification and quantification of pesticide contaminants in food samples. Triclosan is a well-known and common biocide agent against bacteria as well as fungi,²⁰⁵ while bisphenol analogues are used in packaging and lining.²⁰⁶ Morgan et al.²⁰⁶ employed GC–MS to monitor the levels of triclosan and five bisphenol analogues (B, F, P, S, and Z) in 776 adult solid food samples. More than 80% of the samples were contaminated with at least one target phenol. Based on the frequencies, 59% of samples were contaminated with triclosan followed by 32% bisphenol S, and 28% bisphenol Z. The maximum concentration for triclosan was 394 ng/g.

Not only emerging toxins but also the availability of efficient and portable systems have become necessary prerequisites. Some of the recent advancements have shown the availability of portable MS systems for detection and monitoring. Maragos¹³¹ has evaluated the potential of portable MS (APCI-MS) for the detection of T-2 toxin mycotoxin in contaminated cereal grains, wheat, and maize by APCI-MS. The sample was extracted with acetonitrile+water (84:16, v/v) followed by drying and reconstitution in ammonium formate. The MS system contains a linear ion trap mass analyzer to avoid the need of an external supply of gas or air. The device and developed method were able to detect T-2 toxin above 0.2 mg/kg from soft white and hard red wheat, and yellow dent maize. The method was more efficient and hence able to lower-down the detection limit from >0.9 mg/kg. In a similar line, FB and its isoforms were detected in maize with a portable mass spectrometer. For the detection, samples were extracted with aqueous methanol followed by

Table 7. Detection of Emerging Toxins from Food and Water Samples by MS

Name	Sample	Method	Operating parameters	Outcome	Ref
Cypermethrin	Baby food	liquid chromatography coupled to quadrupole Orbitrap mass spectrometry	Gradient elution; negative ionization mode; capillary temperature 300 °C and heater 305 °C	Detection concentration in baby food 10.3 µg kg ⁻¹	208
Parabens	Surface water	UHPLC–MS/MS	LC-18 column; column temp 40 °C; gradient elution; capillary voltage –3.0 kV	Limit of detection 0.04 ng L ⁻¹ and Limit of quantification 0.82 ng L ⁻¹	209
Bisphenol	Water	GS–MS/MS	Temperature transfer line 250 °C; ion source 230 °C and quadrupole 150 °C. solvent delay 4.5 min; electron ionization (EI) mode (70 eV)	Recovery 81.8% –96.1%; limit of detection was 0.2 ng L ⁻¹	210
Parabens	water	GS–MS/MS	Temperature transfer line 250 °C; ion source 230 °C and quadrupole 150 °C. solvent delay 4.5 min; electron ionization (EI) mode (70 eV)	Recovery 81.8% –96.1%; limit of detection was 0.2 ng L ⁻¹	210
Triclosan	water	GS–MS/MS	Temperature transfer line 250 °C; ion source 230 °C and quadrupole 150 °C. solvent delay 4.5 min; electron ionization (EI) mode (70 eV)	Recovery 81.8% –96.1%; limit of detection was 0.2 ng L ⁻¹	210
Neonicotinoids	vegetables	QuEChERS-Portable MS	PDESI as ion source; ultrapure helium (≥99.999%) as carrier gas; inlet temperature 200 °C; molecular pump speed was 1375 Hz	Limit of detection 2.0 ng g ⁻¹ recovery 82.2% –109.7%	211
Carbamates	Vegetables	QuEChERS-Portable MS	PDESI as ion source; ultrapure helium (≥99.999%) as carrier gas; inlet temperature 200 °C; molecular pump speed was 1375 Hz	Limit of detection 2.0 ng g ⁻¹ recovery 82.2% –109.7%	211
Phenyl Pyrazole	Vegetables	QuEChERS-Portable MS	PDESI as ion source; ultrapure helium (≥99.999%) as carrier gas; inlet temperature 200 °C; molecular pump speed was 1375 Hz	Limit of detection 2.0 ng g ⁻¹ recovery 82.2% –109.7%	211
Pesticides	Fruits and vegetables	QuEChERS-LC–MS	Sciex QTRAP 5500 triple quadrupole MS; positive electrospray ionization; ion source temperature 550 °C	24 pesticides detected distinctly	212
Pesticides	Milk	LC-LTQ/Orbitrap Mass Spectrometry	Separation with reverse phase C18 column; positive ionization mode; spray voltage 4 kV; auxiliary gas flow rate 10 arbitrary units; tube lens 90 V, capillary temperature 320 °C	Limit of detection 0.2–8.1 µg kg ⁻¹ and quantification 0.61–24.8 µg kg ⁻¹	213

cleaning up in the immunoaffinity column. Ultimately, cleaned samples were successfully analyzed with the portable MS with detection limits of 0.15 (B1), 0.19 (B2/B3), and 0.28 (total FB) mg/kg maize. The method has quantification limits of 0.33, 0.59, and 0.74 mg/kg and recoveries of 93.6% to 108.6%. Wichert et al.²⁰⁷ have also reported such kind of advancements to detect proteinaceous toxins (912.5–66.5 kDa) from plants as well as microorganisms origin using paper spray-MS (PS-MS) with wipe samples of bench, glass, leaves, flooring, etc., and validated with biological toxin simulant for *Staphylococcal* enterotoxin B. Carbon sputtered porous polyethylene dominated conventional chromatography paper, carbon nanotube-coated paper, and polyethylene for paper spray. The method was able to distinguish the protein toxin simulant efficiently with a good signal-to-noise ratio.

5. DETECTION OF FOOD FRAUD AND FOOD ADULTERATION

Food adulteration and fraud have become a common practice nowadays to gain more profit. To avoid detection by current available analytical methods, new adulteration and fraud practices are becoming advanced and sophisticated making detection one of the biggest challenges for society.²¹⁴ The tracing of fraud in food products via chemical analysis has become more complex especially due to the emergence of new and unknown adulterants.⁷⁴ MS has become an indistinguishable part of testing and food authentication analysis due to its potential to trace chemical compounds based on their chemical fingerprints or chemical profiles.²¹⁵ Adulterants are used as ingredients and additives in food products for economic gain for the seller at the cost of the health of consumers. The use of bulking agents such as sulfated polysaccharides is very common in minced meat and is a fraudulent practice that is very difficult to detect. Kosek et al.²¹⁶ have evaluated rapid evaporative MS (REIMS) to detect adulterants in sausages and burgers prepared from chopped pork and chicken meat. The technique was able to detect the adulterants efficiently with 2.5% as the threshold concentration for protein additives (carrageenan) can be detected at 1% concentration. The major advantage of the system is the quick detection of adulterants in samples, which aids in preventing any major health issue from the consumption of adulterated food products. The problem becomes more serious when consumers are infants or minors. Milk is one of the common products that is supposed to provide optimum nutrients including lipids and proteins. Piras et al.²¹⁷ employed atmospheric pressure matrix-assisted laser desorption/ionization with a Q-TOF mass analyzer, which generated charged proteinaceous as well as lipids/metabolites ions to find the possible fraud with milk. The common adulteration in milk is the intermixing of milk from different sources, and the current methods were able to identify milk from camel, cow, goat, and sheep with 100% accuracy. The goat milk was analyzed for the presence of cow milk as an adulterant, and it was detected even at a lower concentration of 5% with sensitivity and specificity of 92.5% and 94.5%, respectively. The major outcome of the work is its time consumption for sample analysis, i.e., 10 s per unadulterated sample to prepare profile. The method creates differences between protein and lipid molecules in terms of the number of charged moieties as lipids are single charged, while protein moieties have multiple charges.

Integration of artificial intelligence and neural network models has further improved the demand for efficiency of MS systems. Nichani et al.²¹⁸ have developed a method for the

differential identification of spelt and wheat. Nontargeted LC–MS/MS along with convolutional neural network (CNN) models was developed. The employed neural network was able to learn patterns by itself and discriminated between spelt and wheat efficiently. For external validation of the model, artificial mixed spectra of spelt bread and flour, 11 untypical spelt, and six old wheat cultivars (which were not part of model training) were analyzed. The model was able to identify the nonconventional cultivars of wheat and spelt with a *D* value of 0.57.²¹⁸

Not only food products but also active pharma formulations and ingredients are under the threat of adulteration and fraud. One such example is where MS is used to develop a model to differentiate between wild and cultivated *Cordyceps sinensis* harvests. Elemental analysis coupled with stable isotope ratio MS (EA-IRMS) and GasBench II coupled to isotope ratio MS (GB-IRMS) was used with orthogonal partial least-squares discriminant analysis (OPLS-DA) to identify the significant and unique markers of stable isotope ratios in both samples. Analysis identified three stable isotope markers, i.e., $\delta^2\text{H}$, $\delta^{18}\text{O}$, and $\delta^{15}\text{N}$, and their concentrations can be used to identify fresh *C. sinensis* samples as well as differentiating between wild and cultivated *C. sinensis*. $\delta^2\text{H}$ values reduced sequentially in fresh samples based on respective origins. $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ have shown differential patterns as lower $\delta^{18}\text{O}$ and higher $\delta^{15}\text{N}$ represented cultivated samples, while higher $\delta^{18}\text{O}$ and lower $\delta^{15}\text{N}$ represented wild samples. The analyzed cultivated and wild samples have $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ of 18.99%, 4.80%, and 25.72%, 2.29%, respectively.²¹⁹

6. CURRENT CHALLENGES AND FUTURE PROSPECTS

The methods currently in practice for toxin detection are quite advanced, comprising a variety of direct and indirect analytical techniques.²²⁰ However, there are still a few challenges that make laboratory detection difficult, and toxins may sometimes go undetected.²²¹ The following are the significant challenges for toxin analysis that are commonly encountered.

6.1. Representative Sample. Microbial toxins are rarely evenly distributed in foodstuff and edibles.²²² As in the case of mycotoxins, in some regions of the victuals, called “mycotoxin pockets”, concentrations may be extremely high, whereas the rest of the material may be free from contamination. The materials are distributed more heterogeneously for products with larger particle sizes, such as nuts and figs. This necessitates representative sampling that accounts for the random distribution of the “hotspots” to provide an accurate view of the degree of contamination in the specimen. This can be performed by taking a large number of small, incremental samples from various locations distributed throughout the lot.²²³ The selection of incremental samples from the bulk is crucial to give all morsel particles a chance of being selected, thus reducing the statistical bias. Besides the number of samples, sample type and its processing are equally important as in some cases surface swab is sufficient, while others need extraction followed by processing like digestion of target sample as reported in the case of protein-based toxins. Kalb et al. employed endopep-MS to identify botulinum A, B, E, and F from food samples by optimizing the fingerprint peptides generated.¹⁴⁹ Hence, detection technology must be optimized to compete with emerging challenges.

6.2. Sample Preparation. Sample preparation is a highly complex process, with various pitfalls. Every step encountered introduces a level of variability that aggregates and contributes to total variability within single analytical data. It has been generally

observed that nearly 1/3 of the variability is attributed to sample preparation. On the other hand, a much smaller amount of variation is contributed by the analytical method being employed. Even with the best analytical equipment, sample preparation is critical.^{224,225} By adhering to the key factors of sample preparation (size reduction, sampling size, and uniformity), the root-mean-square value can be kept significantly within 5–10%, increasing the prediction result's accuracy.

6.3. Low Concentration. Even at low concentration levels, toxins can be highly toxic.²²⁶ Different classes of toxins have different levels of toxic effects on the target. Glycoalkaloids (potato) and isoflavones (clover) have shown low toxicity, linamarin (cassava) and coniin (hemlock) are somewhat toxic, while ricin (castor beans) and cyanotoxin and saxitoxin (blue-green algae) are extremely toxic.²²⁷ Hence, the tolerable concentration range is also varied in the same proportion. The Food Safety and Standards Authority of India (FSSAI) has set limit values of 15 $\mu\text{g}/\text{kg}$ in cereal products, pulses, and nuts, and 30 $\mu\text{g}/\text{kg}$ in spices, whereas, for milk, the allowable range is considerably low (0.5 $\mu\text{g}/\text{kg}$).²²⁸ The values are more stringent for the US FDA and European Union, nearly 1/3 of the FSSAI-approved figures. The analysis method needs to be extremely precise and sensitive to detect such low concentrations. The low levels of toxin concentration, often not detected, are hugely responsible for reduced production efficiency and increased susceptibility to various diseases.²²⁹

6.4. Complex Matrix. Generally, the toxin matrices found in food are fairly complex and pose a major challenge for laboratories. For example, heterogeneous matrices such as spices contain numerous interfering substances, which makes it extremely difficult to detect the toxins precisely. Also, food may not be necessarily safe, even if well-known toxins are not detected during analysis. These compounds may still be present in conjugated form in masked or bound form.²³⁰ These modified toxins are derived from plants by conjugation and have their chemical structures altered, making the analysis more complex. Also, even though a large number of different toxins including mycotoxins, bacterial toxins, phytotoxins, etc. exist, only a few are characterized and regulated by law.² In reality, several toxins may be present simultaneously, and it may be challenging to find/pinpoint specific analytes in the food particles.

6.5. Portability of Instrument. The normal trend in sample analysis is like the collection of samples from the site and transfer to the lab for further analysis. However, such a protocol seems obsolete when we need to characterize the sample rapidly. In such cases, a portable and handy instrument with easily transferable and ready-to-use techniques is required. Some of the recent research has shown the pay for portable MS-based detection systems for the food analysis and characterization of toxins like PS-MS.²⁰⁷ However, more advancement needed with the miniaturization of instruments with rapid detection and high accuracy is required.

6.6. Cost. A variety of factors contribute to the cost of toxin analysis. They include facilities, sophisticated analytical instruments, reagents, and logistics. Additionally, the number of tests needed for representative sampling to negate the misreporting of toxins in bulk material also contributes to the testing cost. Although effective sampling is one of the most critical factors in mycotoxin analysis, it is the costliest, and surprisingly, innovations in this area have not taken place rapidly.²³¹ Reducing the time and cost requirements through representative sampling while increasing accuracy is the need of the hour. A

balance needs to be achieved between the cost per sample and the number of runs essential to generate the most confident results that will ensure lower downtime with precise results.²³²

6.7. Multiple Toxins Contamination. Naturally, all the food materials are prone to be contaminated with multiple toxins due to the presence of diverse microbial contaminants simultaneously. The method should be effective and efficient to detect all of the toxins together even in minute quantity. A MS-based approach made it possible to allow the detection of diverse range of toxins. Lattanzio et al.²³³ have reported the detection of multiple aflatoxins including B1, B2, G1, G2, ochratoxin A, fumonisins B1 and B2, deoxynivalenol, zearalenone, T-2, and HT-2 toxins simultaneously from maize via LC-MS. In a similar line, Cheng et al.²³⁴ have also reported the detection of 15 toxic alkaloids from vegetables and meat samples using double layer pipet tip magnetic dispersive solid phase extraction method that used polyamidoamine-functionalized magnetic carbon nanotubes. Extracted samples were characterized with ultrafast liquid chromatography-tandem quadrupole mass spectrometry (UFLC-MS/MS) coupled with DPT-MSPE method. The system has shown high recovery efficiency with toxin recovery of up to 125% for meat as well as vegetable dishes. In food samples, fungal contamination is a common phenomenon; hence, it is the prime target for most of the work. Wang et al.²³⁵ and Nualkaw et al.²³⁶ have employed UPLC-MS for the detection of mycotoxins from animal feed like dairy product, poultry, and animal feed. Lee et al.¹⁸⁴ reported the presence of okadaic acid, dinophysistoxin-1, dinophysistoxin-2, and dinophysistoxin-3 in raw as well as cooked mussels using LC-tandem mass spectrometry. The method has shown detection and quantification limit of 0.2–5.1 $\mu\text{g}/\text{kg}$ with accuracy and precision of 80.5–109.8% and 0.9–20.1%, respectively. Albero et al.²³⁷ have also identified mycotoxins in aquaculture feed by LC-MS/MS. The method employed ultrasound-assisted extraction followed by LC-MS/MS, which identified 15 mycotoxins together that also included enniatins (EENB and ENNB1), beauvericin, and fumonisin B2.

In some cases, the toxins are present in modified or bounded form (masked form), which hinder its detection by conventional methods. As mentioned by Berthiller et al.,²³⁸ mycotoxins have high probability of masking due to food processing which changes its structural as well as behavior and obstruct the appearance of common characteristics of respective toxins. Masked mycotoxins have been detected in extractable conjugated and nonextractable varieties (usually to unavailability of toxins in extracted samples, bound mycotoxins are not accessible for analysis and need chemical or enzymatic treatment prior detection).²³⁸

Fiby et al.²³⁹ reported the presence of *Fusarium* mycotoxin like deoxynivalenol in native as well as masked form (DON-3-glucoside "D3G", 3-acetyl-DON "3ADON", or 15-acetyl-DON "15ADON") in cereals. The presence was detected with a stable-isotope dilution liquid chromatography-tandem mass spectrometry-based approach that relies on labeling of toxins enzymatic byproducts with ¹³C. The method efficiently detected D3G (76–98%), DON (86–103%), 15ADON (68–100%), and 3ADON (63–96%).²³⁹ Zhang et al.²⁴⁰ employed ultrahigh-performance liquid chromatography-HRMS to detect 82 mycotoxins and categorize into 8 classes by Python program developed with "Fragmentation pattern screener (FPScreener)" and nontarget screening rules. Pascari et al.²⁴¹ combined QuEChERS with liquid chromatography-triple quadrupole mass spectrometry for the detection of ZEN from

oat flour. For the identification, oat and wheat flours were treated with amylolytic enzymes (α -amylase and amyloglucosidase), similar to the one used in the cereal-based baby food production process that reduced the β -zearalenol (β -ZEL) and β -ZEL-14-sulfate by 40% within 90 min and allowed the specific detection of ZEN-sulfate derivatives from cereals.

7. FUTURE PERSPECTIVE

The past few decades have witnessed significant advancements in analytical techniques and technologies to detect and manage the problem of mycotoxin analysis. The global mycotoxin testing market is estimated to grow at a cumulative annual growth rate (CAGR) of 7.8%, which corresponds to a market of \$1.4 billion by 2026.²⁴² As the number indicates, the accelerated need for food safety has significantly accelerated the mycotoxin testing market. Needless to say, effective sampling, method performance, cost, and rapid detection are of paramount importance for successful mycotoxin testing. However, inexpensive and fast tests with low precision would lead to misclassification through inaccurate results and impact the business decisions of the producers. In light of this, the existing techniques must be refined/tuned for higher precision while active R&D is pursued to develop new methods that can detect multiple toxins simultaneously with high sensitivity (at regulatory levels) with minimal cost and runtime. One option could be developing miniaturized MS, which could be used in a fashion similar to the screening devices used for COVID detection at airports and train stations. The spectrometers are generally efficient and reliable and allow for rapid detection and solid sample detection through thermal absorption. One such example is mycotoxin detection in milled wheat by an MS developed by M/s BaySpec Inc.²⁴³ The portable device could detect even ppm (1.4) levels of the contaminant within less than a minute's time.

Recent studies have highlighted that mass-sensitive microarray (MSMA)-based biosensors could be a promising tool for rapidly detecting mycotoxin material.²⁴⁴ The device consists of a mass-sensitive transducer based on solidly mounted resonator (SMR) technology with a specific integrated circuit. This technology allows small molecular weight toxins (up to 3 toxins simultaneously) to be rapidly detected with high sensitivity (for a single sample) within less than 10 min using mycotoxin analysis as a model example. The authors argued that with further upgrade, many more analytes (~32) could be detected on the devices from a single sample, and with automation, the analysis was performed and printed without the requirement of any operator. The device could be an ideal system for multiplex analysis of mycotoxins; however, significant improvements must be made before it can be considered for field deployment in the analysis of real-time samples.

AUTHOR INFORMATION

Corresponding Author

Kumar Saurav – *Laboratory of Algal Biotechnology-Centre Algatech, Institute of Microbiology of the Czech Academy of Sciences, Třeboň 379 01, Czech Republic*; orcid.org/0000-0002-7084-4204; Email: sauravverma17@gmail.com, saurav@alga.cz

Authors

Vishal Ahuja – *University Institute of Biotechnology, Chandigarh University, Mohali, Punjab 140413, India*; *University Centre for Research & Development, Chandigarh*

University, Mohali, Punjab 140413, India; orcid.org/0000-0001-8982-4978

Amanpreet Singh – *Department of Chemistry, University Institute of Science, Chandigarh University, Mohali, Punjab 140413, India*

Debarati Paul – *Amity Institute of Biotechnology, Noida, Uttar Pradesh 201313, India*

Diptarka Dasgupta – *Material Resource Efficiency Division, CSIR-Indian Institute of Petroleum, Dehradun 248005, India*; orcid.org/0000-0001-9300-8279

Petra Urajová – *Laboratory of Algal Biotechnology-Centre Algatech, Institute of Microbiology of the Czech Academy of Sciences, Třeboň 379 01, Czech Republic*

Sounak Ghosh – *Laboratory of Algal Biotechnology-Centre Algatech, Institute of Microbiology of the Czech Academy of Sciences, Třeboň 379 01, Czech Republic*

Roshani Singh – *Laboratory of Algal Biotechnology-Centre Algatech, Institute of Microbiology of the Czech Academy of Sciences, Třeboň 379 01, Czech Republic*

Gobardhan Sahoo – *Laboratory of Algal Biotechnology-Centre Algatech, Institute of Microbiology of the Czech Academy of Sciences, Třeboň 379 01, Czech Republic*

Daniela Ewe – *Laboratory of Algal Biotechnology-Centre Algatech, Institute of Microbiology of the Czech Academy of Sciences, Třeboň 379 01, Czech Republic*

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.chemrestox.3c00241>

Author Contributions

V.A., and K.S.: conceptualization, investigation, data analysis, original draft preparation, and reviewing. A.S., D.P., D.D., S.S., R.S., G.S., D.E., and P.U.: original draft preparation, editing, and reviewing. K.S.: supervision, data analysis, and reviewing. All authors have read and agreed to the final version of the manuscript. CRediT: **Vishal Ahuja** conceptualization, data curation, formal analysis, methodology, software, validation, visualization, writing-original draft, writing-review & editing; **Amanpreet Singh** writing-original draft; **Debarati Paul** writing-original draft; **Diptarka Dasgupta** writing-original draft; **Petra Urajová** software, validation, visualization, writing-review & editing; **Sounak Ghosh** writing-review & editing; **Roshani Singh** writing-review & editing; **Gobardhan Sahoo** formal analysis, visualization; **Daniela Ewe** formal analysis, validation, writing-review & editing; **Kumar Saurav** conceptualization, formal analysis, investigation, project administration, supervision, writing-original draft, writing-review & editing.

Notes

The authors declare no competing financial interest.

Biographies



Dr. Vishal Ahuja has his Ph.D. in Biotechnology from Himachal Pradesh University, Shimla, India. He has his post doc in CSIR-Indian Institute of Petroleum Dehradun India on an FTT project. Currently he is working as assistant professor of research and development (UIBT/UCRD) Chandigarh University, Punjab, India. He has published more than 30 research and review articles. His research interest is in biomass biotransformation, production of bioactive compounds, and energy products from sustainable and renewable resources. His area of work is closely aligned with current trends in biotechnology advancements in fields such as biofuels, pharmaceuticals, and environmental sustainability.



Dr. Amanpreet Singh received his Ph.D. degree from the Department of Chemistry, Indian Institute of Technology, Ropar in 2023. He earned his master's degree in organic chemistry from the Department of Chemistry, Panjab University Chandigarh in 2017. Currently, he is an assistant professor in the Department of Chemistry, University Institute of Science, Chandigarh University, Mohali. He has published his research in well reputed international journals. His research interest lies in supramolecular organic chemistry, material chemistry, and development of carbon-based nanomaterials for sensing and catalysis. His research mainly focuses on fabrication of chemo sensors for environmental applications and sustainable development.




Dr. Debarati Paul, currently Associate Professor in Amity Institute of Biotechnology, Amity University, has completed her Ph.D. from the Institute of Microbial Technology-CSIR (Chandigarh) and post doc from EAWAG (Switzerland) and Mississippi State University (USA) for about 5 years. Her main area of research includes genomics, environmental biotechnology, antimicrobial resistance, and biodegradation. She has published over 70 papers including research articles, reviews, conference proceedings and posters, and book chapters. She has received the prestigious Indo US Science and Technology Forum award and DST travel grants, etc. She also trains students for skill development in Quality Control Chemist Microbiology, since 2019.



Dr. Diptarka Dasgupta is working as principal scientist in Material Resource Efficiency Division of CSIR-Indian Institute of Petroleum Dehradun Uttarakhand India. He has completed his Ph.D. from AcSIR-Indian Institute of Petroleum. He has expertise in microbial genomics and biotransformation of waste to industrial products. He has published more than 40 research and review articles. He is also experienced in analytical techniques for characterization of biomolecules.



Dr. Petra Urajová received her Ph.D. from University of South Bohemia, Department of Agricultural Chemistry. She has published over 25 research articles in well reputed international journals. Her research is mainly focused on chemistry of natural products, specifically algae and cyanobacteria and their secondary metabolites. Among her skills belong mainly HPLC and HPLC–HRMS techniques.



Sounak Ghosh is currently working on his final year dissertation project at Laboratory of Algal Biotechnology, Institute of Microbiology of Czech Academy of Sciences, Czech Republic. He is an ongoing integrated master's student in biotechnology enrolled at Vellore Institute of Technology, Vellore, India. He has worked on multiple projects including Bioremediation and microbiology. His current research work is focused on genome mining of secondary metabolites, and natural product isolation from various symbiotic cyanobacteria.



Roshani Singh is currently working on her final year dissertation project at Laboratory of Algal Biotechnology, Institute of Microbiology of Czech Academy of Sciences, Czech Republic. She is an ongoing integrated master's student in biotechnology enrolled at Vellore Institute of Technology, Vellore, India. She has worked on various projects on bioinformatics, bioremediation, and molecular biology. She is currently focused on exploration of novel cyanobacterial metabolites by direct pathway cloning.



Dr. Gobardhan Sahoo received his Ph.D. from CSIR-National Institute of Oceanography under the aegis of Academy of Scientific and Innovative Research (AcSIR), India in Science category. Further, he has pursued his postdoctoral studies at Department of Marine Microbiology, University of Haifa, Israel. Currently, he is pursuing his postdoctoral study at Laboratory of Algal Biotechnology, Institute of Microbiology of Czech Academy of Sciences, Czech Republic. He has published more than 11 research papers in national and international journals. His research is mainly focused on the ecology of epibiotic diatoms and the biocommunication system in barnacles.



Dr. Daniela Ewe received her Ph.D. from Department of Biology, University of Konstanz, Germany in 2015. She published over 15 research articles in well reputed international journals. Further, she has pursued her postdoctoral studies at Laboratory of Algal Biotechnology, Institute of Microbiology of Czech Academy of Sciences, Czech Republic and CeBiTec – Center for Biotechnology, Bielefeld University, Germany. She has vast experience in the field of microalgae physiology and its metabolism.



Dr. Kumar Saurav received his Ph.D. from Vellore Institute of Technology, India followed by two postdoctoral studies at South China Sea Institute of Oceanology, Chinese Academy of Sciences and Department of Marine Microbiology, University of Haifa, Israel. Currently, he is working as junior group leader at Laboratory of Algal Biotechnology, Institute of Microbiology of Czech Academy of Sciences, Czech Republic. He has published over 50 research articles in well reputed international journals. He has vast experience in the isolation of secondary metabolites from microorganisms and their biological activity determination. He is well versed in the analytical instruments like LC–MS, HPLC, and HRMS.

REFERENCES

- (1) WHO Estimates of the Global Burden of Foodborne Diseases: Foodborne Disease Burden Epidemiology Reference Group 2007–2015; World Health Organization, 2015.
- (2) Alshannaq, A.; Yu, J.-H. Occurrence, Toxicity, and Analysis of Major Mycotoxins in Food. *Int. J. Environ. Res. Public Health* **2017**, *14* (6), 632.
- (3) Rasheed, T.; Bilal, M.; Nabeel, F.; Adeel, M.; Iqbal, H. M. N. Environmentally-Related Contaminants of High Concern: Potential Sources and Analytical Modalities for Detection, Quantification, and Treatment. *Environ. Int.* **2019**, *122*, 52–66.
- (4) Rather, I. A.; Koh, W. Y.; Paek, W. K.; Lim, J. The Sources of Chemical Contaminants in Food and Their Health Implications. *Front Pharmacol* **2017**, *8*, 830.
- (5) Fletcher, M. T.; Netzel, G. Food Safety and Natural Toxins. *Toxins* **2020**, *12* (4), 236.
- (6) Dey, P.; Kundu, A.; Kumar, A.; Gupta, M.; Lee, B. M.; Bhakta, T.; Dash, S.; Kim, H. S. Analysis of Alkaloids (Indole Alkaloids, Isoquinoline Alkaloids, Tropane Alkaloids). *Recent Advances in Natural Products Analysis* **2020**, 505–567.
- (7) Picardo, M.; Núñez, O.; Farré, M. Suspect and Target Screening of Natural Toxins in the Ter River Catchment Area in NE Spain and Prioritisation by Their Toxicity. *Toxins* **2020**, *12* (12), 752.
- (8) Malachová, A.; Stránská, M.; Václavíková, M.; Elliott, C. T.; Black, C.; Meneely, J.; Hajšlová, J.; Ezekiel, C. N.; Schuhmacher, R.; Krška, R. Advanced LC-MS-Based Methods to Study the Co-Occurrence and Metabolization of Multiple Mycotoxins in Cereals and Cereal-Based Food. *Anal Bioanal Chem.* **2018**, *410* (3), 801–825.
- (9) Krška, R.; de Nijs, M.; McNerney, O.; Pichler, M.; Gilbert, J.; Edwards, S.; Suman, M.; Magan, N.; Rossi, V.; van der Fels-Klerx, H. J.; Bagi, F.; Poschmaier, B.; Sulyok, M.; Berthiller, F.; van Egmond, H. P. Safe Food and Feed through an Integrated Toolbox for Mycotoxin Management: The MyToolBox Approach. *World Mycotoxin Journal* **2016**, *9* (4), 487–495.
- (10) Turner, N. W.; Subrahmanyam, S.; Piletsky, S. A. Analytical Methods for Determination of Mycotoxins: A Review. *Anal. Chim. Acta* **2009**, *632* (2), 168–180.
- (11) Verma, M.; Chaudhary, M.; Singh, A.; Kaur, N.; Singh, N. Naphthalimide-Gold-Based Nanocomposite for the Ratiometric Detection of Okadaic Acid in Shellfish. *J. Mater. Chem. B* **2020**, *8* (36), 8405–8413.
- (12) Fayyaz, K.; Nawaz, A.; Olaimat, A. N.; Akram, K.; Farooq, U.; Fatima, M.; Siddiqui, S. A.; Rana, I. S.; Mahnoor, M.; Shahbaz, H. M. Microbial Toxins in Fermented Foods: Health Implications and Analytical Techniques for Detection. *Journal of Food and Drug Analysis* **2022**, *30* (4), 523–537.
- (13) Abeyesiriwardena, N. M.; Gascoigne, S. J. L.; Anandappa, A. Algal Bloom Expansion Increases Cyanotoxin Risk in Food. *Yale J. Biol. Med.* **2018**, *91* (2), 129–142.
- (14) Mtewa, A. G.; Egbuna, C.; Ngwira, K. J.; Lampiao, F.; Shah, U.; Mtewa, T. K. Classification of Phytotoxins and Their Mechanisms of Action. In *Poisonous Plants and Phytochemicals in Drug Discovery*; John Wiley & Sons, Ltd, 2020; pp 19–30. DOI: 10.1002/9781119650034.ch2.
- (15) Campo, J.; Picó, Y. 17 - Emerging Contaminants and Toxins. In *Chemical Analysis of Food (Second ed.)*; Pico, Y., Ed.; Academic Press, 2020; pp 729–758. DOI: 10.1016/B978-0-12-813266-1.00017-6.
- (16) Dhakal, A.; Hashmi, M. F.; Sbar, E. Aflatoxin Toxicity. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, 2023.
- (17) Jastrzebski, M. K.; Kaczor, A. A.; Wróbel, T. M. Methods of Lysergic Acid Synthesis-The Key Ergot Alkaloid. *Molecules* **2022**, *27* (21), 7322.
- (18) Maragos, C. M.; Barnett, K.; Morgan, L.; Vaughan, M. M.; Sieve, K. K. Measurement of Fumonisin in Maize Using a Portable Mass Spectrometer. *Toxins* **2022**, *14* (8), 523.
- (19) Wang, L.; Wang, Q.; Wang, S.; Cai, R.; Yuan, Y.; Yue, T.; Wang, Z. Bio-Control on the Contamination of Ochratoxin A in Food: Current Research and Future Prospects. *Curr. Res. Food Sci.* **2022**, *5*, 1539–1549.
- (20) Bacha, S. A. S.; Li, Y.; Nie, J.; Xu, G.; Han, L.; Farooq, S. Comprehensive Review on Patulin and Alternaria Toxins in Fruit and Derived Products. *Frontiers in Plant Science* **2023**, *14*. DOI: 10.3389/fpls.2023.1139757
- (21) Jing, S.; Liu, C.; Zheng, J.; Dong, Z.; Guo, N. Toxicity of Zearalenone and Its Nutritional Intervention by Natural Products. *Food Funct.* **2022**, *13* (20), 10374–10400.
- (22) Puváča, N.; Avantaggiato, G.; Merkuri, J.; Vuković, G.; Bursić, V.; Cara, M. Occurrence and Determination of Alternaria Mycotoxins Alternariol, Alternariol Monomethyl Ether, and Tentoxin in Wheat Grains by QuEChERS Method. *Toxins (Basel)* **2022**, *14* (11), 791.
- (23) Charla, R.; Patil, P. P.; Patil, V. S.; Bhandare, V. V.; Karoshi, V.; Balaganur, V.; Joshi, R. K.; Harish, D. R.; Roy, S. Anti-Cholera Toxin Activity of Selected Polyphenols from Careya Arborea, Punica Granatum, and Psidium Guajava. *Frontiers in Cellular and Infection Microbiology* **2023**, *13*. DOI: 10.3389/fcimb.2023.1106293
- (24) Pomputius, W. F.; Kilgore, S. H.; Schlievert, P. M. Probable Enterotoxin-Associated Toxic Shock Syndrome Caused by Staphylococcus Epidermidis. *BMC Pediatr* **2023**, *23* (1), 108.
- (25) Nada, H. G.; El-Tahan, A. S.; El-Didamony, G.; Askora, A. Detection of Multidrug-Resistant Shiga Toxin-Producing Escherichia Coli in Some Food Products and Cattle Faeces in Al-Sharkia, Egypt: One Health Menace. *BMC Microbiology* **2023**, *23* (1), 127.
- (26) Zhang, L.; Lv, Q.; Zheng, Y.; Gao, S.; Huang, W.; Liu, P.; Kong, D.; Wang, Y.; Yu, Y.; Jiang, Y.; Jiang, H. Rapid and Sensitive Detection of Botulinum Toxin Type A in Complex Sample Matrices by AlphaLISA. *Frontiers in Public Health* **2022**, *10*. DOI: 10.3389/fpubh.2022.987517
- (27) Yang, S.; Wang, Y.; Liu, Y.; Jia, K.; Zhang, Z.; Dong, Q. Cereulide and Emetic Bacillus Cereus: Characterizations, Impacts and Public Precautions. *Foods* **2023**, *12* (4), 833.
- (28) Wang, T.; Fernandes, S. P. S.; Araújo, J.; Li, X.; Salonen, L. M.; Espiña, B. A Carboxyl-Functionalized Covalent Organic Polymer for the Efficient Adsorption of Saxitoxin. *J. Hazard Mater.* **2023**, *452*, 131247.
- (29) Panlilio, J. M.; Hammar, K. M.; Aluru, N.; Hahn, M. E. Developmental Exposure to Domoic Acid Targets Reticulospinal Neurons and Leads to Aberrant Myelination in the Spinal Cord. *Sci. Rep* **2023**, *13* (1), 2587.
- (30) Tebben, J.; Zurhelle, C.; Tubaro, A.; Samdal, I. A.; Krock, B.; Kilcoyne, J.; Sosa, S.; Trainer, V. L.; Deeds, J. R.; Tillmann, U. Structure and Toxicity of AZA-59, an Azaspiracid Shellfish Poisoning Toxin Produced by Azadinium Poporum (Dinophyceae). *Harmful Algae* **2023**, *124*, 102388.
- (31) Brammer-Robbins, E.; Costa, K. A.; Bowden, J. A.; Martyniuk, C. J.; Larkin, I. V.; Denslow, N. D. Putative High-Level Toxicity Pathways Based on Evidence of Brevetoxin Immunotoxicity in Marine Fauna. *Aquat Toxicol* **2022**, *252*, 106298.
- (32) Qin, Y.; Li, J.; Kuang, J.; Shen, S.; Jiang, J.; Zhang, Z.; Zhao, C.; Zhou, X.; Huang, B.; Han, B. Sensitive Time-Resolved Fluoroimmunoassay for the Quantitative Detection of Okadaic Acid. *Frontiers in Marine Science* **2022**, *9*. DOI: 10.3389/fmars.2022.961751
- (33) Bolarinwa, I. F.; Oke, M. O.; Olaniyan, S. A.; Ajala, A. S.; Bolarinwa, I. F.; Oke, M. O.; Olaniyan, S. A.; Ajala, A. S. A Review of

Cyanogenic Glycosides in Edible Plants. In *Toxicology - New Aspects to This Scientific Conundrum*; IntechOpen, 2016. DOI: 10.5772/64886.

- (34) Melough, M. M.; Lee, S. G.; Cho, E.; Kim, K.; Provatas, A. A.; Perkins, C.; Park, M. K.; Qureshi, A.; Chun, O. K. Identification and Quantitation of Furocoumarins in Popularly Consumed Foods in the U.S. Using QuEChERS Extraction Coupled with UPLC-MS/MS Analysis. *J. Agric. Food Chem.* **2017**, *65* (24), 5049–5055.
- (35) Wu, J. S.; Clauson-Kaas, F.; Lindqvist, D. N.; Rasmussen, L. H.; Strobel, B. W.; Hansen, H. C. B. Does the Natural Carcinogen Ptaquiloside Degrade Readily in Groundwater? *Environmental Sciences Europe* **2021**, *33* (1), 24.
- (36) Stegelmeier, B. L.; Colegate, S. M.; Brown, A. W. Dehydropyrrolizidine Alkaloid Toxicity, Cytotoxicity, and Carcinogenicity. *Toxins (Basel)* **2016**, *8* (12), 356.
- (37) Girardi, C.; Odore, R. Pharmacological Treatments and Risks for the Food Chain. *Vet Res. Commun.* **2008**, *32*, S11–18.
- (38) Kralj Cigic, I.; Prosen, H. An Overview of Conventional and Emerging Analytical Methods for the Determination of Mycotoxins. *Int. J. Mol. Sci.* **2009**, *10* (1), 62–115.
- (39) Kole, P. L.; Venkatesh, G.; Kotecha, J.; Sheshala, R. Recent Advances in Sample Preparation Techniques for Effective Bioanalytical Methods. *Biomed Chromatogr* **2011**, *25* (1–2), 199–217.
- (40) Qiu, N.; Sun, D.; Zhou, S.; Li, J.; Zhao, Y.; Wu, Y. Rapid and Sensitive UHPLC-MS/MS Methods for Dietary Sample Analysis of 43 Mycotoxins in China Total Diet Study. *Journal of Advanced Research* **2022**, *39*, 15–47.
- (41) De Girolamo, A.; Lippolis, V.; Pascale, M. Overview of Recent Liquid Chromatography Mass Spectrometry-Based Methods for Natural Toxins Detection in Food Products. *Toxins (Basel)* **2022**, *14* (5), 328.
- (42) von Bargen, C.; Brockmeyer, J.; Humpf, H.-U. Meat Authentication: A New HPLC-MS/MS Based Method for the Fast and Sensitive Detection of Horse and Pork in Highly Processed Food. *J. Agric. Food Chem.* **2014**, *62* (39), 9428–9435.
- (43) Liu, Z.; Zhang, M.; Chen, P.; Harnly, J. M.; Sun, J. Mass Spectrometry-Based Nontargeted and Targeted Analytical Approaches in Fingerprinting and Metabolomics of Food and Agricultural Research. *J. Agric. Food Chem.* **2022**, *70* (36), 11138–11153.
- (44) Monbaliu, S.; Van Poucke, C.; Detavernier, C.; Dumoulin, F.; Van De Velde, M.; Schoeters, E.; Van Dyck, S.; Averkieva, O.; Van Peteghem, C.; De Saeger, S. Occurrence of Mycotoxins in Feed as Analyzed by a Multi-Mycotoxin LC-MS/MS Method. *J. Agric. Food Chem.* **2010**, *58* (1), 66–71.
- (45) Kerian, K. S.; Jarmusch, A. K.; Cooks, R. G. Touch Spray Mass Spectrometry for In Situ Analysis of Complex Samples. *Analyst* **2014**, *139* (11), 2714–2720.
- (46) Feider, C. L.; Krieger, A.; DeHoog, R. J.; Eberlin, L. S. Ambient Ionization Mass Spectrometry: Recent Developments and Applications. *Anal. Chem.* **2019**, *91* (7), 4266–4290.
- (47) Hernández-Mesa, M.; Le Bizec, B.; Dervilly, G. Metabolomics in Chemical Risk Analysis - A Review. *Anal. Chim. Acta* **2021**, *1154*, 338298.
- (48) Selamat, J.; Rozani, N. A. A.; Murugesu, S. Application of the Metabolomics Approach in Food Authentication. *Molecules* **2021**, *26* (24), 7565.
- (49) Vilariño, N.; Louzao, M. C.; Fraga, M.; Rodríguez, L. P.; Botana, L. M. Innovative Detection Methods for Aquatic Algal Toxins and Their Presence in the Food Chain. *Anal Bioanal Chem.* **2013**, *405* (24), 7719–7732.
- (50) Law, J. W.-F.; Ab Mutalib, N.-S.; Chan, K.-G.; Lee, L.-H. Rapid Methods for the Detection of Foodborne Bacterial Pathogens: Principles, Applications, Advantages and Limitations. *Front Microbiol* **2015**, *5*, 770.
- (51) Liew, W.-P.-P.; Sabran, M.-R. Recent Advances in Immunoassay-Based Mycotoxin Analysis and Toxicogenomic Technologies. *J. Food Drug Anal* **2022**, *30* (4), 549–561.
- (52) Quilliam, M. A.; Thomas, K.; Wright, J. L. Analysis of Domoic Acid in Shellfish by Thin-Layer Chromatography. *Nat. Toxins* **1998**, *6* (3–4), 147–152.
- (53) Rodríguez, I.; Vieytes, M. R.; Alfonso, A. Analytical Challenges for Regulated Marine Toxins. Detection Methods. *Current Opinion in Food Science* **2017**, *18*, 29–36.
- (54) Jangampalli Adi, P.; Matcha, B. Analysis of Aflatoxin B1 in Contaminated Feed, Media, and Serum Samples of *Cyprinus Carpio* L. by High-Performance Liquid Chromatography. *Food Quality and Safety* **2018**, *2* (4), 199–204.
- (55) Siracusa, M.; Bacchicocchi, S.; Dubbini, A.; Campacci, D.; Tavoloni, T.; Stramenga, A.; Ciriaci, M.; Dall'Ara, S.; Piersanti, A. A High Throughput Screening HPLC-FLD Method for Paralytic Shellfish Toxins (PSTs) Enabling Effective Official Control. *Molecules* **2022**, *27* (15), 4702.
- (56) Chandreleka, T.; Abinayaa, S.; Rathour, A.; Mahalakshmi, A.; Heema, C. Detection of Aflatoxin in Food Products Using UV Fluorescence Spectroscopy. *IJITEE* **2020**, *9* (5), 1068–1071.
- (57) Singh, A. K.; Lakshmi, G. B. V. S.; Dhiman, T. K.; Kaushik, A.; Solanki, P. R. Bio-Active Free Direct Optical Sensing of Aflatoxin B1 and Ochratoxin A Using a Manganese Oxide Nano-System. *Frontiers in Nanotechnology* **2021**, *2*. DOI: 10.3389/fnano.2020.621681
- (58) Ray, S. Sensory Properties of Foods and Their Measurement Methods. In *Techniques to Measure Food Safety and Quality: Microbial, Chemical, and Sensory*; Khan, M. S., Shafiur Rahman, M., Eds.; Springer International Publishing: Cham, 2021; pp 345–381. DOI: 10.1007/978-3-030-68636-9_15.
- (59) Liu, Y.; Qiao, F.; Xu, L.; Wang, R.; Jiang, W.; Xu, Z. Fast Detection of Diarrhetic Shellfish Poisoning Toxins in Mussels Using NIR Spectroscopy and Improved Twin Support Vector Machines. *Frontiers in Marine Science* **2022**, *9*. DOI: 10.3389/fmars.2022.907378
- (60) Baden, D. G.; Melinek, R.; Sechet, V.; Trainer, V. L.; Schultz, D. R.; Rein, K. S.; Tomas, C. R.; Delgado, J.; Hale, L. Modified Immunoassays for Polyether Toxins: Implications of Biological Matrixes, Metabolic States, and Epitope Recognition. *Journal of AOAC INTERNATIONAL* **1995**, *78* (2), 499–507.
- (61) Yasumoto, T.; Murata, M.; Oshima, Y.; Sano, M.; Matsumoto, G. K.; Clardy, J. Diarrhetic Shellfish Toxins. *Tetrahedron* **1985**, *41* (6), 1019–1025.
- (62) Tubaro, A.; Florio, C.; Luxich, E.; Sosa, S.; Della Loggia, R.; Yasumoto, T. A Protein Phosphatase 2A Inhibition Assay for a Fast and Sensitive Assessment of Okadaic Acid Contamination in Mussels. *Toxicon* **1996**, *34* (7), 743–752.
- (63) Eberhart, B.-T. L.; Moore, L. K.; Harrington, N.; Adams, N. G.; Borchert, J.; Trainer, V. L. Screening Tests for the Rapid Detection of Diarrhetic Shellfish Toxins in Washington State. *Mar Drugs* **2013**, *11* (10), 3718–3734.
- (64) Kim, H. U.; Goepfert, J. M. Enumeration and Identification of *Bacillus Cereus* in Foods. I. 24-h Presumptive Test Medium. *Appl. Microbiol.* **1971**, *22* (4), 581–587.
- (65) Lund, T.; Granum, P. E. Comparison of Biological Effect of the Two Different Enterotoxin Complexes Isolated from Three Different Strains of *Bacillus Cereus*. *Microbiology (Reading)* **1997**, *143* (10), 3329–3336.
- (66) Hasan, M. R.; Pulingam, T.; Appaturi, J. N.; Zifruddin, A. N.; Teh, S. J.; Lim, T. W.; Ibrahim, F.; Leo, B. F.; Thong, K. L. Carbon Nanotube-Based Aptasensor for Sensitive Electrochemical Detection of Whole-Cell *Salmonella*. *Anal. Biochem.* **2018**, *554*, 34–43.
- (67) Tevell Åberg, A.; Björnstad, K.; Hedeland, M. Mass Spectrometric Detection of Protein-Based Toxins. *Biosecurity and Bioterrorism: Biodefense Strategy, Practice, and Science* **2013**, *11* (S1), S215–S226.
- (68) Silva, C. J. Food Forensics: Using Mass Spectrometry To Detect Foodborne Protein Contaminants, as Exemplified by Shiga Toxin Variants and Prion Strains. *J. Agric. Food Chem.* **2018**, *66* (32), 8435–8450.
- (69) Rodríguez-Carrasco, Y.; Moltó, J. C.; Mañes, J.; Berrada, H. Development of a GC-MS/MS Strategy to Determine 15 Mycotoxins and Metabolites in Human Urine. *Talanta* **2014**, *128*, 125–131.
- (70) Jadhav, S. R.; Shah, R. M.; Karpe, A. V.; Morrison, P. D.; Kouremenos, K.; Beale, D. J.; Palombo, E. A. Detection of Foodborne

Pathogens Using Proteomics and Metabolomics-Based Approaches. *Frontiers in Microbiology* **2018**, *9*. DOI: 10.3389/fmicb.2018.03132

(71) Xu, M.-L.; Gao, Y.; Wang, X.; Han, X. X.; Zhao, B. Comprehensive Strategy for Sample Preparation for the Analysis of Food Contaminants and Residues by GC-MS/MS: A Review of Recent Research Trends. *Foods* **2021**, *10* (10), 2473.

(72) Kalb, S. R.; Boyer, A. E.; Barr, J. R. Mass Spectrometric Detection of Bacterial Protein Toxins and Their Enzymatic Activity. *Toxins* **2015**, *7* (9), 3497–3511.

(73) Pavlovic, M.; Huber, I.; Konrad, R.; Busch, U. Application of MALDI-TOF MS for the Identification of Food Borne Bacteria. *Open Microbiol J.* **2013**, *7*, 135–141.

(74) Zamboni, C. MALDI-TOF Mass Spectrometry Applications for Food Fraud Detection. *Applied Sciences* **2021**, *11* (8), 3374.

(75) Xing, L. Determination of Toxic Elements in Food by ICP-MS Using AOAC Method 2015.01. *Spectroscopy* **2022**, *37* (10), 7–12.

(76) Ammann, A. A. Inductively Coupled Plasma Mass Spectrometry (ICP MS): A Versatile Tool. *Journal of Mass Spectrometry* **2007**, *42* (4), 419–427.

(77) te Brinke, E.; Arrizabalaga-Larrañaga, A.; Blokland, M. H. Insights of Ion Mobility Spectrometry and Its Application on Food Safety and Authenticity: A Review. *Anal. Chim. Acta* **2022**, *1222*, 340039.

(78) Hernández-Mesa, M.; Ropartz, D.; García-Campaña, A. M.; Rogniaux, H.; Dervilly-Pinel, G.; Le Bizec, B. Ion Mobility Spectrometry in Food Analysis: Principles, Current Applications and Future Trends. *Molecules* **2019**, *24* (15), 2706.

(79) Büyükköröglü, G.; Dora, D. D.; Özdemir, F.; Hızal, C. Chapter 15 - Techniques for Protein Analysis. In *Omics Technologies and Bio-Engineering*; Barh, D.; Azevedo, V., Eds.; Academic Press, 2018; pp 317–351. DOI: 10.1016/B978-0-12-804659-3.00015-4.

(80) Domínguez, I.; Frenich, A. G.; Romero-González, R. Mass Spectrometry Approaches to Ensure Food Safety. *Analytical Methods* **2020**, *12* (9), 1148–1162.

(81) Mazarakioti, E. C.; Zotos, A.; Thomatou, A. A.; Kontogeorgos, A.; Patakas, A.; Ladavos, A. Inductively Coupled Plasma-Mass Spectrometry (ICP-MS), a Useful Tool in Authenticity of Agricultural Products' and Foods' Origin. *Foods* **2022**, *11* (22), 3705.

(82) Jull, A. J. T.; Burr, G. S. 5.20 - Accelerator Mass Spectrometry. In *Treatise on Geochemistry (Second ed.)*; Holland, H. D., Turekian, K. K., Eds.; Elsevier: Oxford, 2014; pp 375–383. DOI: 10.1016/B978-0-08-095975-7.01429-7.

(83) Wacker, J. F.; Eiden, G. C.; Lehn, S. A.; Koppelaar, D. W. Accelerator Mass Spectrometry (AMS). In *Encyclopedia of Spectroscopy and Spectrometry (Third ed.)*; Lindon, J. C., Tranter, G. E., Koppelaar, D. W., Eds.; Academic Press: Oxford, 2017; pp 15–17. DOI: 10.1016/B978-0-12-803224-4.00043-1.

(84) Metz, T. O.; Zhang, Q.; Page, J. S.; Shen, Y.; Callister, S. J.; Jacobs, J. M.; Smith, R. D. The Future of Liquid Chromatography-Mass Spectrometry (LC-MS) in Metabolic Profiling and Metabolomic Studies for Biomarker Discovery. *Biomark Med.* **2007**, *1* (1), 159–185.

(85) Alves, T. O.; D'Almeida, C. T. S.; Scherf, K. A.; Ferreira, M. S. L. Modern Approaches in the Identification and Quantification of Immunogenic Peptides in Cereals by LC-MS/MS. *Frontiers in Plant Science* **2019**, *10*. DOI: 10.3389/fpls.2019.01470

(86) Amirav, A.; Fialkov, A.; Margolin Eren, K.; Neumark, B.; Elkabets, O.; Tsizin, S.; Gordin, A.; Alon, T. *Gas Chromatography-Mass Spectrometry (GC-MS) with Cold Electron Ionization (EI): Bridging the Gap Between GC-MS and LC-MS.* **2020**, *18* (4), 5–15.

(87) Gilbert, J. R.; McCaskill, D.; Fishman, V. N.; Brzak, K.; Markham, D.; Bartels, M. J.; Winniford, B.; Bano Mohsin, S.; Godbey, J.; Akinbo, O.; Lewer, P. Chapter 17 - Industrial Applications of High-Resolution GC/MS. In *Comprehensive Analytical Chemistry*; Ferrer, I., Thurman, E. M., Eds.; Advanced Techniques in Gas Chromatography-Mass Spectrometry (GC-MS-MS and GC-TOF-MS) for Environmental Chemistry; Elsevier, 2013; Vol. 61, pp 403–429. DOI: 10.1016/B978-0-444-62623-3.00017-4.

(88) Vivekanandan-Giri, A.; Byun, J.; Pennathur, S. Chapter Five - Quantitative Analysis of Amino Acid Oxidation Markers by Tandem

Mass Spectrometry. In *Methods in Enzymology*; Conn, P. M., Ed.; The Unfolded Protein Response and Cellular Stress, Part C; Academic Press, 2011; Vol. 491, pp 73–89. DOI: 10.1016/B978-0-12-385928-0.00005-5.

(89) Geer Wallace, M. A.; McCord, J. P. Chapter 16 - High-Resolution Mass Spectrometry. In *Breathborne Biomarkers and the Human Volatilome (Second ed.)*; Beauchamp, J., Davis, C., Pleil, J., Eds.; Elsevier: Boston, 2020; pp 253–270. DOI: 10.1016/B978-0-12-819967-1.00016-5.

(90) Rychert, J. Benefits and Limitations of MALDI-TOF Mass Spectrometry for the Identification of Microorganisms. *Journal of Infectiology and Epidemiology* **2019**, *2* (4), 1.

(91) Wilschefski, S. C.; Baxter, M. R. Inductively Coupled Plasma Mass Spectrometry: Introduction to Analytical Aspects. *Clin Biochem Rev.* **2019**, *40* (3), 115–133.

(92) Vorderwülbecke, S.; Cleverley, S.; Weinberger, S. R.; Wiesner, A. Protein Quantification by the SELDI-TOF-MS-Based ProteinChip® System. *Nat. Methods* **2005**, *2* (5), 393–395.

(93) Poon, T. C. Opportunities and Limitations of SELDI-TOF-MS in Biomedical Research: Practical Advices. *Expert Review of Proteomics* **2007**, *4* (1), 51–65.

(94) Neagu, A.-N.; Jayathirtha, M.; Baxter, E.; Donnelly, M.; Petre, B. A.; Darie, C. C. Applications of Tandem Mass Spectrometry (MS/MS) in Protein Analysis for Biomedical Research. *Molecules* **2022**, *27* (8), 2411.

(95) Mittal, R. D. Tandem Mass Spectroscopy in Diagnosis and Clinical Research. *Indian J. Clin Biochem* **2015**, *30* (2), 121–123.

(96) Anđjelković, U.; Šrajer Gajdošik, M.; Gašo-Sokač, D.; Martinović, T.; Josić, D. Foodomics and Food Safety: Where We Are. *Food Technol. Biotechnol* **2017**, *55* (3), 290–307.

(97) Wu, P.; Wu, X.; Huang, Q.; Yu, Q.; et al. Mass Spectrometry-Based Multimodal Approaches for the Identification and Quantification Analysis of Microplastics in Food Matrix. *Front. Nutr.* **2023**, *10* (April), 1–7.

(98) Spanjer, M. C. *Mass Spectrometry in Multi-Mycotoxin and Fungal Spore Analysis* **2011**, 90.

(99) Yang, S.; Zhu, X.; Wang, J.; Jin, X.; Liu, Y.; Qian, F.; Zhang, S.; Chen, J. Combustion of Hazardous Biological Waste Derived from the Fermentation of Antibiotics Using TG-FTIR and Py-GC/MS Techniques. *Bioresour. Technol.* **2015**, *193* (x), 156–163.

(100) Zhou, Q.; Xing, A.; Zhao, K. Simultaneous Determination of Nickel, Cobalt and Mercury Ions in Water Samples by Solid Phase Extraction Using Multiwalled Carbon Nanotubes as Adsorbent after Chelating with Sodium Diethyldithiocarbamate Prior to High Performance Liquid Chromatography. *Journal of Chromatography A* **2014**, *1360*, 76–81.

(101) El-Sayed, R. A.; Jebur, A. B.; Kang, W.; El-Demerdash, F. M. An Overview on the Major Mycotoxins in Food Products: Characteristics, Toxicity, and Analysis. *Journal of Future Foods* **2022**, *2* (2), 91–102.

(102) Yu, S.; Jia, B.; Li, K.; Zhou, H.; Lai, W.; Tang, Y.; Yan, Z.; Sun, W.; Liu, N.; Yu, D.; Wu, A. Pre-Warning of Abiotic Factors in Maize Required for Potential Contamination of Fusarium Mycotoxins via Response Surface Analysis. *Food Control* **2021**, *121*, 107570.

(103) Bennett, J. W.; Klich, M. *Mycotoxins.* *Clin Microbiol Rev.* **2003**, *16* (3), 497–516.

(104) Mohammadi Shad, Z.; Venkitasamy, C. Mycotoxins as Food and Feed Contaminant: Effect on Health and Economy and Their Management. In *Fungal Resources for Sustainable Economy: Current Status and Future Perspectives*; Singh, I., Rajpal, V. R., Navi, S. S., Eds.; Springer Nature: Singapore, 2023; pp 531–563. DOI: 10.1007/978-981-19-9103-5_20.

(105) Kumar, P.; Mahato, D. K.; Kamle, M.; Mohanta, T. K.; Kang, S. G. Aflatoxins: A Global Concern for Food Safety, Human Health and Their Management. *Frontiers in Microbiology* **2017**, *7*. DOI: 10.3389/fmicb.2016.02170

(106) Tirmenstein, M. A.; Mangipudy, R. Aflatoxin. In *Encyclopedia of Toxicology (Third ed.)*; Wexler, P., Ed.; Academic Press: Oxford, 2014; pp 104–106. DOI: 10.1016/B978-0-12-386454-3.00224-4.

- (107) Pinto, A. C. S. M.; De Pierri, C. R.; Evangelista, A. G.; Gomes, A. S. d. L. P. B.; Luciano, F. B. Deoxynivalenol: Toxicology, Degradation by Bacteria, and Phylogenetic Analysis. *Toxins* **2022**, *14* (2), 90.
- (108) Cope, R. B. Chapter 75 - Trichothecenes. In *Veterinary Toxicology (Third ed.)*; Gupta, R. C., Ed.; Academic Press, 2018; pp 1043–1053. DOI: 10.1016/B978-0-12-811410-0.00075-1.
- (109) Kamle, M.; Mahato, D. K.; Devi, S.; Lee, K. E.; Kang, S. G.; Kumar, P. Fumonisin: Impact on Agriculture, Food, and Human Health and Their Management Strategies. *Toxins (Basel)* **2019**, *11* (6), 328.
- (110) Braun, M. S.; Wink, M. Exposure, Occurrence, and Chemistry of Fumonisin and Their Cryptic Derivatives. *Comprehensive Reviews in Food Science and Food Safety* **2018**, *17* (3), 769–791.
- (111) Damiani, T.; Righetti, L.; Suman, M.; Galaverna, G.; Dall'Asta, C. Analytical Issue Related to Fumonisin: A Matter of Sample Comminution? *Food Control* **2019**, *95*, 1–5.
- (112) Ropejko, K.; Twaruzek, M. Zearalenone and Its Metabolites—General Overview, Occurrence, and Toxicity. *Toxins (Basel)* **2021**, *13* (1), 35.
- (113) Zhang, G.-L.; Feng, Y.-L.; Song, J.-L.; Zhou, X.-S. Zearalenone: A Mycotoxin With Different Toxic Effect in Domestic and Laboratory Animals' Granulosa Cells. *Frontiers in Genetics* **2018**, *9*. DOI: 10.3389/fgene.2018.00667
- (114) Ben Salah-Abbès, J.; Belgacem, H.; Ezzdini, K.; Abdel-Wahhab, M. A.; Abbès, S. Zearalenone Nephrotoxicity: DNA Fragmentation, Apoptotic Gene Expression and Oxidative Stress Protected by *Lactobacillus Plantarum* MON03. *Toxicon* **2020**, *175*, 28–35.
- (115) Gromadzka, K.; Waskiewicz, A.; Chelkowski, J.; Golinski, P. Zearalenone and Its Metabolites: Occurrence, Detection, Toxicity and Guidelines. *World Mycotoxin Journal* **2008**, *1* (2), 209–220.
- (116) Mally, A.; Solfrizzo, M.; Degen, G. H. Biomonitoring of the Mycotoxin Zearalenone: Current State-of-the Art and Application to Human Exposure Assessment. *Arch. Toxicol.* **2016**, *90* (6), 1281–1292.
- (117) Reddy, P.; Hemsworth, J.; Guthridge, K. M.; Vinh, A.; Vassiliadis, S.; Ezernieks, V.; Spangenberg, G. C.; Rochfort, S. J. Ergot Alkaloid Mycotoxins: Physiological Effects, Metabolism and Distribution of the Residual Toxin in Mice. *Sci. Rep* **2020**, *10* (1), 9714.
- (118) Schiff, P. L. Ergot and Its Alkaloids. *Am. J. Pharm. Educ* **2006**, *70* (5), 98.
- (119) Evans, T. J. Chapter 67 - Endocrine Disruptors. In *Reproductive and Developmental Toxicology*; Gupta, R. C., Ed.; Academic Press: San Diego, 2011; pp 873–891. DOI: 10.1016/B978-0-12-382032-7.10067-0.
- (120) Bui-Klimke, T. R.; Wu, F. Ochratoxin A and Human Health Risk: A Review of the Evidence. *Crit. Rev. Food Sci. Nutr* **2015**, *55* (13), 1860–1869.
- (121) Al-Jaal, B.; Salama, S.; Al-Qasbi, N.; Jaganjac, M. Mycotoxin Contamination of Food and Feed in the Gulf Cooperation Council Countries and Its Detection. *Toxicon* **2019**, *171*, 43–50.
- (122) European Union Commission. *Commission Regulation (EC) No 401/2006 of 23 February 2006 Laying down the Methods of Sampling and Analysis for the Official Control of the Levels of Mycotoxins in Foodstuffs*; EU, 2006; Vol. 401/2006, p 12. <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CONSLEG:2006R0401:20100313:EN:PDF>.
- (123) European Union Commission. *COMMISSION REGULATION (EC) No 1881/2006 of 19 December 2006 Setting Maximum Levels for Certain Contaminants in Foodstuffs*; EU, 2006; Vol. 1881/2006. <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2006:364:0005:0024:EN:PDF>.
- (124) European Union Commission. *Commission Regulation (EC) No 1881/2006 of 19 December 2006 Setting Maximum Levels for Certain Contaminants in Foodstuffs (Text with EEA Relevance)*; EU, 2006; Vol. 364. <http://data.europa.eu/eli/reg/2006/1881/oj/eng> (accessed 05-28-2023).
- (125) European Union Commission. *2013/165/EU: Commission Recommendation of 27 March 2013 on the Presence of T-2 and HT-2 Toxin in Cereals and Cereal Products Text with EEA Relevance*; EU, 2013; Vol. OJ L 91, 3.4.2013, pp 12–15. <https://eur-lex.europa.eu/eli/reco/2013/165/oj#PP3Contents> (accessed 05-28-2023).
- (126) European Union Commission. *Commission Recommendation of 17 August 2006 on the Presence of Deoxynivalenol, Zearalenone, Ochratoxin A, T-2 and HT-2 and Fumonisin in Products Intended for Animal Feeding (Text with EEA Relevance) (2006/576/EC)*; EU, 2016. <http://data.europa.eu/eli/reco/2006/576/2016-08-02/eng> (accessed 05-28-2023).
- (127) European Union Commission. *Directive 2002/32/EC of the European Parliament and of the Council of 7 May 2002 on Undesirable Substances in Animal Feed*; EU, 2019. <http://data.europa.eu/eli/dir/2002/32/2019-11-28/eng> (accessed 05-28-2023).
- (128) Mbughuni, M. M.; Jannetto, P. J.; Langman, L. J. Mass Spectrometry Applications for Toxicology. *EJIFCC* **2016**, *27* (4), 272–287.
- (129) Areo, O. M.; Abafe, O. A.; Gbashi, S.; Njobeh, P. B. Detection of Multi-Mycotoxins in Rooibos and Other Consumed Teas in South Africa by a Modified QuEChERS Method and Ultra-High Performance Liquid Chromatography Tandem Mass Spectrometry. *Food Control* **2023**, *143*, 109255.
- (130) You, Y.; Hu, Q.; Liu, N.; Xu, C.; Lu, S.; Xu, T.; Mao, X. Metabolite Analysis of *Alternaria* Mycotoxins by LC-MS/MS and Multiple Tools. *Molecules* **2023**, *28* (7), 3258.
- (131) Maragos, C. M. Detection of T-2 Toxin in Wheat and Maize with a Portable Mass Spectrometer. *Toxins (Basel)* **2023**, *15* (3), 222.
- (132) Ji, X.; Deng, T.; Xiao, Y.; Jin, C.; Lyu, W.; Wang, W.; Tang, B.; Wu, Z.; Yang, H. Evaluation of *Alternaria* Toxins in Fruits, Vegetables and Their Derivatives Marketed in China Using a QuEChERS Method Coupled with Ultra-High Performance Liquid Chromatography-Tandem Mass Spectrometry: Analytical Methods and Occurrence. *Food Control* **2023**, *147*, 109563.
- (133) Leite, M.; Freitas, A.; Barbosa, J.; Ramos, F. Mycotoxins in Raw Bovine Milk: UHPLC-QTrap-MS/MS Method as a Biosafety Control Tool. *Toxins* **2023**, *15* (3), 173.
- (134) Chen, M.; Liu, X.; Yang, S.; Chen, Z.; Di, B.; Liu, W.; Yan, H. HPLC-MS/MS Method for the Simultaneous Determination of Aflatoxins in Blood: Toxicokinetics of Aflatoxin B1 and Aflatoxin M1 in Rats. *Journal of Analytical Science and Technology* **2022**, *13* (1), 27.
- (135) Tang, H.; Han, W.; Fei, S.; Li, Y.; Huang, J.; Dong, M.; Wang, L.; Wang, W.; Zhang, Y. Development of Acid Hydrolysis-Based UPLC-MS/MS Method for Determination of *Alternaria* Toxins and Its Application in the Occurrence Assessment in Solanaceous Vegetables and Their Products. *Toxins* **2023**, *15* (3), 201.
- (136) Cai, C.; Liu, Y.; Tang, X.; Zhang, W.; Zhang, Q.; Li, P. Development of a Toxin-Free Competitive Immunoassay for Aflatoxin M1 Based on a Nanobody as Surrogate Calibrator. *LWT* **2023**, *182*, 114829.
- (137) Prakasham, K.; Gurrani, S.; Shiea, J.-T.; Wu, M.-T.; Wu, C.-F.; Ku, Y.-J.; Tsai, T.-Y.; Hua, H.-T.; Lin, Y.-J.; Huang, P.-C.; Andaluri, G.; Ponnusamy, V. K. Rapid Identification and Analysis of Ochratoxin-A in Food and Agricultural Soil Samples Using a Novel Semi-Automated In-Syringe Based Fast Mycotoxin Extraction (FaMEx) Technique Coupled with UHPLC-MS/MS. *Molecules* **2023**, *28* (3), 1442.
- (138) Prusova, N.; Behner, A.; Dzuman, Z.; Hajslova, J.; Stranska, M. Conjugated Type A Trichothecenes in Oat-Based Products: Occurrence Data and Estimation of the Related Risk. *Food Control* **2023**, *143*, 109281.
- (139) Scheibenzuber, S.; Dick, F.; Asam, S.; Rychlik, M. Analysis of 13 *Alternaria* Mycotoxins Including Modified Forms in Beer. *Mycotoxin Res.* **2021**, *37* (2), 149–159.
- (140) Rodríguez-Carrasco, Y.; Mañes, J.; Berrada, H.; Juan, C. Development and Validation of a LC-ESI-MS/MS Method for the Determination of *Alternaria* Toxins Alternariol, Alternariol Methyl-Ether and Tentoxin in Tomato and Tomato-Based Products. *Toxins (Basel)* **2016**, *8* (11), 328.
- (141) Bielecki, J. Emerging Food Pathogens and Bacterial Toxins. *Acta Microbiol Pol* **2003**, *52*, 17–22.
- (142) Hernández-Cortez, C.; Palma-Martínez, I.; UrielGonzalez-Avila, L.; Guerrero-Mandujano, A.; Castro-Escarpulli, R. C. S. G.; Hernández-Cortez, C.; Palma-Martínez, I.; UrielGonzalez-Avila, L.; Guerrero-Mandujano, A.; Castro-Escarpulli, R. C. S. G. Food Poisoning

- Caused by Bacteria (Food Toxins). In *Poisoning - From Specific Toxic Agents to Novel Rapid and Simplified Techniques for Analysis*; IntechOpen, 2017. DOI: 10.5772/intechopen.69953.
- (143) Romero Hurtado, S.; Iregui, C. El Lipopolisacárido. *Revista de Medicina Veterinaria* **2010**, *1* (19), 37–45.
- (144) Ramachandran, G. Gram-Positive and Gram-Negative Bacterial Toxins in Sepsis. *Virulence* **2014**, *5* (1), 213–218.
- (145) Foran, P.; Lawrence, G. W.; Shone, C. C.; Foster, K. A.; Dolly, J. O. Botulinum Neurotoxin C1 Cleaves Both Syntaxin and SNAP-25 in Intact and Permeabilized Chromaffin Cells: Correlation with Its Blockade of Catecholamine Release. *Biochemistry* **1996**, *35* (8), 2630–2636.
- (146) Schiavo, G. G.; Benfenati, F.; Poulain, B.; Rossetto, O.; de Laureto, P. P.; DasGupta, B. R.; Montecucco, C. Tetanus and Botulinum-B Neurotoxins Block Neurotransmitter Release by Proteolytic Cleavage of Synaptobrevin. *Nature* **1992**, *359* (6398), 832–835.
- (147) Kuo, S.-R.; Willingham, M. C.; Bour, S. H.; Andreas, E. A.; Park, S. K.; Jackson, C.; Duesbery, N. S.; Leppla, S. H.; Tang, W.-J.; Frankel, A. E. Anthrax Toxin-Induced Shock in Rats Is Associated with Pulmonary Edema and Hemorrhage. *Microbial Pathogenesis* **2008**, *44* (6), 467–472.
- (148) Duracova, M.; Klimentova, J.; Myslivcova Fucikova, A.; Zidkova, L.; Sheshko, V.; Rehulkova, H.; Dresler, J.; Krocova, Z. Targeted Mass Spectrometry Analysis of Clostridium Perfringens Toxins. *Toxins (Basel)* **2019**, *11* (3), 177.
- (149) Kalb, S. R.; Krilich, J. C.; Dykes, J. K.; Lúquez, C.; Maslanka, S. E.; Barr, J. R. Detection of Botulinum Toxins A, B, E, and F in Foods by Endopep-MS. *J. Agric. Food Chem.* **2015**, *63* (4), 1133–1141.
- (150) Rosen, O.; Feldberg, L.; Yamin, T. S.; Dor, E.; Barnea, A.; Weissberg, A.; Zichel, R. Development of a Multiplex Endopep-MS Assay for Simultaneous Detection of Botulinum Toxins A, B and E. *Sci. Rep.* **2017**, *7* (1), 14859.
- (151) Drigo, I.; Tonon, E.; Pascoletti, S.; Anniballi, F.; Kalb, S. R.; Bano, L. Detection of Active BoNT/C and D by EndoPep-MS Using MALDI Biotyper Instrument and Comparison with the Mouse Test Bioassay. *Toxins* **2021**, *13* (1), 10.
- (152) Wang, H.; Hu, L.; Chang, X.; Hu, Y.; Zhang, Y.; Zhou, P.; Cui, X. Determination of Bacterial Toxin Toxoflavin and Fervenuin in Food and Identification of Their Degradation Products. *Food Chem.* **2023**, *399*, 134010.
- (153) Safi, A. U. R.; Mansour Salih, M.; Rahman, H.; Khattak, B.; El Askary, A.; Hussain Khalifa, E.; Qasim, M. Immunoaffinity-Based Mass Spectrometric Characterization of Immunoreactive Proteins of Salmonella Typhi. *Saudi Journal of Biological Sciences* **2023**, *30* (1), 103502.
- (154) Murray, P. R. What Is New in Clinical Microbiology—Microbial Identification by MALDI-TOF Mass Spectrometry. *J. Mol. Diagn.* **2012**, *14* (5), 419–423.
- (155) Fagerquist, C. K.; Garbus, B. R.; Miller, W. G.; Williams, K. E.; Yee, E.; Bates, A. H.; Boyle, S.; Harden, L. A.; Cooley, M. B.; Mandrell, R. E. Rapid Identification of Protein Biomarkers of Escherichia Coli O157:H7 by Matrix-Assisted Laser Desorption Ionization-Time-of-Flight-Time-of-Flight Mass Spectrometry and Top-down Proteomics. *Anal. Chem.* **2010**, *82* (7), 2717–2725.
- (156) Gagnaire, J.; Dauwalder, O.; Boisset, S.; Khau, D.; Freydiere, A.-M.; Ader, F.; Bes, M.; Lina, G.; Tristan, A.; Reverdy, M.-E.; Marchand, A.; Geissmann, T.; Benito, Y.; Durand, G.; Charrier, J.-P.; Etienne, J.; Welker, M.; Van Belkum, A.; Vandenesch, F. Detection of Staphylococcus Aureus Delta-Toxin Production by Whole-Cell MALDI-TOF Mass Spectrometry. *PLoS One* **2012**, *7* (7), No. e40660.
- (157) Thakur, M. S.; Ragavan, K. V. Biosensors in Food Processing. *J. Food Sci. Technol.* **2013**, *50* (4), 625–641.
- (158) Bragazzi, N. L.; Amicizia, D.; Panatto, D.; Tramalloni, D.; Valle, I.; Gasparini, R. Quartz-Crystal Microbalance (QCM) for Public Health: An Overview of Its Applications. *Adv. Protein Chem. Struct. Biol.* **2015**, *101*, 149–211.
- (159) Lamanna, J.; Scott, E. Y.; Edwards, H. S.; Chamberlain, M. D.; Dryden, M. D. M.; Peng, J.; Mair, B.; Lee, A.; Chan, C.; Sklavounos, A. A.; Heffernan, A.; Abbas, F.; Lam, C.; Olson, M. E.; Moffat, J.; Wheeler, A. R. Digital Microfluidic Isolation of Single Cells for -Omics. *Nat. Commun.* **2020**, *11* (1), 5632.
- (160) Nedelkov, D.; Rasooly, A.; Nelson, R. W. Multitoxin Biosensor-Mass Spectrometry Analysis: A New Approach for Rapid, Real-Time, Sensitive Analysis of Staphylococcal Toxins in Food. *Int. J. Food Microbiol.* **2000**, *60* (1), 1–13.
- (161) Visciano, P.; Schirone, M.; Berti, M.; Milandri, A.; Tofalo, R.; Suzzi, G. Marine Biotoxins: Occurrence, Toxicity, Regulatory Limits and Reference Methods. *Frontiers in Microbiology* **2016**, *7*. DOI: 10.3389/fmicb.2016.01051
- (162) Rambla-Alegre, M.; Miles, C. O.; de la Iglesia, P.; Fernandez-Tejedor, M.; Jacobs, S.; Sioen, I.; Verbeke, W.; Samdal, I. A.; Sandvik, M.; Barbosa, V.; Tediosi, A.; Madorran, E.; Granby, K.; Kotterman, M.; Calis, T.; Diogene, J. Occurrence of Cyclic Imines in European Commercial Seafood and Consumers Risk Assessment. *Environmental Research* **2018**, *161*, 392–398.
- (163) García-Altare, M.; Casanova, A.; Bane, V.; Diogène, J.; Furey, A.; De la Iglesia, P. Confirmation of Pinnatoxins and Spirolides in Shellfish and Passive Samplers from Catalonia (Spain) by Liquid Chromatography Coupled with Triple Quadrupole and High-Resolution Hybrid Tandem Mass Spectrometry. *Marine Drugs* **2014**, *12* (6), 3706–3732.
- (164) Otero, P.; Pérez, S.; Alfonso, A.; Vale, C.; Rodríguez, P.; Gouveia, N. N.; Gouveia, N.; Delgado, J.; Vale, P.; Hiram, M.; Ishihara, Y.; Molgó, J.; Botana, L. M. First Toxin Profile of Ciguateric Fish in Madeira Arquipélago (Europe). *Anal. Chem.* **2010**, *82* (14), 6032–6039.
- (165) Estevez, P.; Castro, D.; Pequeño-Valtierra, A.; Giraldez, J.; Gago-Martinez, A. Emerging Marine Biotoxins in Seafood from European Coasts: Incidence and Analytical Challenges. *Foods* **2019**, *8* (5), 149.
- (166) Burrell, S.; Gunnarsson, T.; Gunnarsson, K.; Clarke, D.; Turner, A. D. First Detection of Paralytic Shellfish Poisoning (PSP) Toxins in Icelandic Mussels (*Mytilus Edulis*): Links to Causative Phytoplankton Species. *Food Control* **2013**, *31* (2), 295–301.
- (167) SCHWARZ, M.; JANDOVÁ, K.; STRUK, I.; MAREŠOVÁ, D.; POKORNÝ, J.; RILJAK, V. Low Dose Domoic Acid Influences Spontaneous Behavior in Adult Rats. *Physiological Research* **2014**, *63*, 369–376.
- (168) Mafra, L. L.; Bricelj, V. M.; Fennel, K. Domoic Acid Uptake and Elimination Kinetics in Oysters and Mussels in Relation to Body Size and Anatomical Distribution of Toxin. *Aquatic Toxicology* **2010**, *100* (1), 17–29.
- (169) Nielsen, L. T.; Hansen, P. J.; Krock, B.; Vismann, B. Accumulation, Transformation and Breakdown of DSP Toxins from the Toxic Dinoflagellate Dinophysis Acuta in Blue Mussels, *Mytilus Edulis*. *Toxicon* **2016**, *117*, 84–93.
- (170) Louzao, M. C.; Vilariño, N.; Vale, C.; Costas, C.; Cao, A.; Raposo-Garcia, S.; Vieytes, M. R.; Botana, L. M. Current Trends and New Challenges in Marine Phycotoxins. *Mar Drugs* **2022**, *20* (3), 198.
- (171) Li, A.; Ma, J.; Cao, J.; McCarron, P. Toxins in Mussels (*Mytilus Galloprovincialis*) Associated with Diarrhetic Shellfish Poisoning Episodes in China. *Toxicon* **2012**, *60* (3), 420–425.
- (172) Sun, P.; Leeson, C.; Zhi, X.; Leng, F.; Pierce, R. H.; Henry, M. S.; Rein, K. S. Characterization of an Epoxide Hydrolase from the Florida Red Tide Dinoflagellate, *Karenia Brevis*. *Phytochemistry* **2016**, *122*, 11–21.
- (173) Traylor, J.; Singhal, M. Ciguatera Toxicity. In *StatPearls*; StatPearls Publishing: Treasure Island (FL), 2023.
- (174) Pottier, I.; Lewis, R. J.; Vernoux, J.-P. Ciguatera Fish Poisoning in the Caribbean Sea and Atlantic Ocean: Reconciling the Multiplicity of Ciguatoxins and Analytical Chemistry Approach for Public Health Safety. *Toxins* **2023**, *15* (7), 453.
- (175) Hidalgo, J.; Liberona, J. L.; Molgó, J.; Jaimovich, E. Pacific Ciguatoxin-1b Effect over Na⁺ and K⁺ Currents, Inositol 1,4,5-Triphosphate Content and Intracellular Ca²⁺ Signals in Cultured Rat Myotubes. *Br. J. Pharmacol.* **2002**, *137* (7), 1055–1062.

- (176) Vijayakumar, G.; Venkatesan, S. A.; Kannan, V. A.; Perumal, S. Chapter 8 - Detection of Food Toxins, Pathogens, and Microorganisms Using Nanotechnology-Based Sensors. In *Nanotechnology Applications for Food Safety and Quality Monitoring*; Sharma, A., Vijayakumar, P. S., Prabhakar, E. P. K., Kumar, R., Eds.; Academic Press, 2023; pp 155–170. DOI: 10.1016/B978-0-323-85791-8.00022-7.
- (177) Karunarathne, A.; Bhalla, A.; Sethi, A.; Perera, U.; Eddleston, M. Importance of Pesticides for Lethal Poisoning in India during 1999 to 2018: A Systematic Review. *BMC Public Health* **2021**, *21*, 1441.
- (178) Blay, P.; Hui, J. P. M.; Chang, J.; Melanson, J. E. Screening for Multiple Classes of Marine Biotoxins by Liquid Chromatography-High-Resolution Mass Spectrometry. *Anal Bioanal Chem.* **2011**, *400* (2), 577–585.
- (179) Bosch-Orea, C.; Sanchís, J.; Farré, M. Analysis of Highly Polar Marine Biotoxins in Seawater by Hydrophilic Interaction Liquid Chromatography Coupled to High Resolution Mass Spectrometry. *MethodsX* **2021**, *8*, 101370.
- (180) Kolrep, F.; Hessel, S.; These, A.; Ehlers, A.; Rein, K.; Lampen, A. Differences in Metabolism of the Marine Biotoxin Okadaic Acid by Human and Rat Cytochrome P450 Monooxygenases. *Arch. Toxicol.* **2016**, *90* (8), 2025–2036.
- (181) Kolrep, F.; Rein, K.; Lampen, A.; Hessel-Pras, S. Metabolism of Okadaic Acid by NADPH-Dependent Enzymes Present in Human or Rat Liver S9 Fractions Results in Different Toxic Effects. *Toxicology in Vitro* **2017**, *42*, 161–170.
- (182) Lefebvre, D.; Blanco-Valle, K.; Hennekinne, J.-A.; Simon, S.; Fenaillé, F.; Becher, F.; Nia, Y. Multiplex Detection of 24 Staphylococcal Enterotoxins in Culture Supernatant Using Liquid Chromatography Coupled to High-Resolution Mass Spectrometry. *Toxins (Basel)* **2022**, *14* (4), 249.
- (183) Barr, J. R.; Moura, H.; Boyer, A. E.; Woolfitt, A. R.; Kalb, S. R.; Pavlopoulos, A.; McWilliams, L. G.; Schmidt, J. G.; Martinez, R. A.; Ashley, D. L. Botulinum Neurotoxin Detection and Differentiation by Mass Spectrometry. *Emerg Infect Dis* **2005**, *11* (10), 1578–1583.
- (184) Lee, S. Y.; Woo, S. Y.; Tian, F.; Park, J. B.; Choi, K.-S.; Chun, H. S. Simultaneous Determination of Okadaic Acid, Dinophysistoxin-1, Dinophysistoxin-2, and Dinophysistoxin-3 Using Liquid Chromatography-Tandem Mass Spectrometry in Raw and Cooked Food Matrices. *Food Control* **2022**, *139*, 109068.
- (185) Beach, D. G.; Zamylyny, L.; MacArthur, M.; Miles, C. O. Liquid Chromatography-High-Resolution Tandem Mass Spectrometry of Anatoxins, Including New Conjugates and Reduction Products. *Anal Bioanal Chem.* **2023**, *415*, 5281.
- (186) Aparicio-Muriana, M. d. M.; Lara, F. J.; Olmo-Iruela, M. D.; García-Campana, A. M. Determination of Multiclass Cyanotoxins in Blue-Green Algae (BGA) Dietary Supplements Using Hydrophilic Interaction Liquid Chromatography-Tandem Mass Spectrometry. *Toxins* **2023**, *15* (2), 127.
- (187) Li, P.; Lu, R.; Zhu, J.; Ke, Q.; Wang, D.; Wang, Y.; Zhang, C.; Lu, X.; Zhao, M.; Wang, Z.; Pan, L.; Zhang, M.; Ying, J.; Li, S. Simultaneous Determination of 13 Paralytic Shellfish Toxins and Tetrodotoxin in Marine Shellfish by Porous Graphitic Carbon Solid-Phase Extraction and Hydrophilic Interaction Chromatography-Tandem Mass Spectrometry. *Food Additives & Contaminants: Part A* **2023**, *40* (1), 147–158.
- (188) Urugo, M. M.; Tringo, T. T. Naturally Occurring Plant Food Toxicants and the Role of Food Processing Methods in Their Detoxification. *Int. J. Food Sci.* **2023**, *2023*, 9947841.
- (189) Colegate, S. M.; Gardner, D. R.; Lee, S. T. Plant-Associated Natural Food Toxins. In *Handbook of Food Chemistry*; Cheung, P. C. K., Mehta, B. M., Eds.; Springer Berlin Heidelberg: Berlin, Heidelberg, 2015; pp 753–783. DOI: 10.1007/978-3-642-36605-5_9.
- (190) Arigò, A.; Rigano, F.; Russo, M.; Trovato, E.; Dugo, P.; Mondello, L. Dietary Intake of Coumarins and Furocoumarins through Citrus Beverages: A Detailed Estimation by a HPLC-MS/MS Method Combined with the Linear Retention Index System. *Foods* **2021**, *10* (7), 1533.
- (191) Rizzo, S.; Celano, R.; Piccinelli, A. L.; Serio, S.; Russo, M.; Rastrelli, L. An Analytical Platform for the Screening and Identification of Pyrrolizidine Alkaloids in Food Matrices with High Risk of Contamination. *Food Chem.* **2023**, *406*, 135058.
- (192) Bessaire, T.; Ernest, M.; Christinat, N.; Carrères, B.; Panchaud, A.; Badoud, F. High Resolution Mass Spectrometry Workflow for the Analysis of Food Contaminants: Application to Plant Toxins, Mycotoxins and Phytoestrogens in Plant-Based Ingredients. *Food Addit Contam Part A Chem. Anal Control Expo Risk Assess* **2021**, *38* (6), 978–996.
- (193) Kalb, S. R.; Schieltz, D. M.; Becher, F.; Astot, C.; Fredriksson, S.-Å.; Barr, J. R. Recommended Mass Spectrometry-Based Strategies to Identify Ricin-Containing Samples. *Toxins (Basel)* **2015**, *7* (12), 4881–4894.
- (194) Liang, X.; Christensen, J. H.; Nielsen, N. J. Enhancing the Power of Liquid Chromatography-Mass Spectrometry for Chemical Fingerprinting of Phytotoxins in the Environment. *Journal of Chromatography A* **2021**, *1642*, 462027.
- (195) Hosu, I. S.; Sobaszek, M.; Ficek, M.; Bogdanowicz, R.; Coffinier, Y. Boron-Doped Carbon Nanowalls for Fast and Direct Detection of Cytochrome C and Ricin by Matrix-Free Laser Desorption/Ionization Mass Spectrometry. *Talanta* **2023**, *252*, 123778.
- (196) Chen, L.; Zhang, Q.; Yi, Z.; Chen, Y.; Xiao, W.; Su, D.; Shi, W. Risk Assessment of (Herbal) Teas Containing Pyrrolizidine Alkaloids (PAs) Based on Margin of Exposure Approach and Relative Potency (REP) Factors. *Foods* **2022**, *11* (19), 2946.
- (197) Park, H.; Cho, Y.; Lee, J.; Lee, K. M.; Kim, H. J.; Lee, J.; Bahn, Y.-S.; Son, J. Evaluation and Monitoring of the Natural Toxin Ptaquiloside in Bracken Fern, Meat, and Dairy Products. *Toxins* **2023**, *15* (3), 231.
- (198) Dosoky, N. S.; Satyal, P.; Setzer, W. N. Authentication of Citrus Spp. Cold-Pressed Essential Oils by Their Oxygenated Heterocyclic Components. *Molecules* **2022**, *27* (19), 6277.
- (199) Makovi, C. M.; Parker, C. H.; Zhang, K. Determination of Amygdalin in Apricot Kernels and Almonds Using LC-MS/MS. *J. AOAC Int.* **2023**, *106* (2), 457–463.
- (200) Rizzo, S.; Celano, R.; Piccinelli, A. L.; Russo, M.; Rastrelli, L. Target Screening Method for the Quantitative Determination of 118 Pyrrolizidine Alkaloids in Food Supplements, Herbal Infusions, Honey and Teas by Liquid Chromatography Coupled to Quadrupole Orbitrap Mass Spectrometry. *Food Chem.* **2023**, *423*, 136306.
- (201) Gruber-Dorninger, C.; Novak, B.; Nagl, V.; Berthiller, F. Emerging Mycotoxins: Beyond Traditionally Determined Food Contaminants. *J. Agric. Food Chem.* **2017**, *65* (33), 7052–7070.
- (202) Thompson, L. A.; Darwish, W. S. Environmental Chemical Contaminants in Food: Review of a Global Problem. *J. Toxicol* **2019**, *2019*, 2345283.
- (203) Sage, A.; Taylor, P.; Stahl-Zeng, J. Novel Methods Using Mass Spectrometry for Food Safety—From Contamination to Nutrition. *Column* **2018**, *14* (5), 13–19.
- (204) Vass, A.; Robles-Molina, J.; Pérez-Ortega, P.; Gilbert-López, B.; Dernovics, M.; Molina-Díaz, A.; García-Reyes, J. F. Study of Different HILIC, Mixed-Mode, and Other Aqueous Normal-Phase Approaches for the Liquid Chromatography/Mass Spectrometry-Based Determination of Challenging Polar Pesticides. *Anal Bioanal Chem.* **2016**, *408* (18), 4857–4869.
- (205) Gopalakrishnan, S.; Ghosh, R.; Renganathan, T.; Pushpavanam, S. Sensitive and Selective Determination of Triclosan Using Visual Spectroscopy. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* **2021**, *254*, 119623.
- (206) Morgan, M. K.; Clifton, M. S. Exposure to Triclosan and Bisphenol Analogues B, F, P, S and Z in Repeated Duplicate-Diet Solid Food Samples of Adults. *Toxics* **2021**, *9* (3), 47.
- (207) Wichert, W. R. A.; Dhummakupt, E. S.; Zhang, C.; Mach, P. M.; Bernhards, R. C.; Glaros, T.; Manicke, N. E. Detection of Protein Toxin Simulants from Contaminated Surfaces by Paper Spray Mass Spectrometry. *J. Am. Soc. Mass Spectrom.* **2019**, *30* (8), 1406–1415.
- (208) Prata, R.; López-Ruiz, R.; Petrarca, M. H.; Teixeira Godoy, H.; Garrido Frenich, A.; Romero-González, R. Targeted and Non-Targeted Analysis of Pesticides and Aflatoxins in Baby Foods by Liquid

Chromatography Coupled to Quadrupole Orbitrap Mass Spectrometry. *Food Control* **2022**, *139*, 109072.

(209) Thuy, Q. C.; Phuong, P. T.; Thien, T. L. T.; Minh, B. Q. Determination of Seven Parabens in Surface Water Samples by UHPLC-MS/MS and Solid-Phase Extraction. *Vietnam Journal of Chemistry* **2022**, *60* (6), 738–743.

(210) Bian, L.; Li, S.; Ge, X.; Wang, M.; Li, K.; Wang, X. Determination of Bisphenols, Triclosan, and Parabens in Bottled Water by Solid-Phase Microextraction Combined with Gas Chromatography-Tandem Mass Spectrometry and Assessment of the Associated Health Risk. *Journal of Food Composition and Analysis* **2023**, *123*, 105548.

(211) Mou, B.; Zuo, C.; Chen, L.; Xie, H.; Zhang, W.; Wang, Q.; Wen, L.; Gan, N. On-Site Simultaneous Determination of Neonicotinoids, Carbamates, and Phenyl Pyrazole Insecticides in Vegetables by QuEChERS Extraction on Nitrogen and Sulfur Co-Doped Carbon Dots and Portable Mass Spectrometry. *Journal of Chromatography A* **2023**, *1689*, 463744.

(212) Zhang, S.; He, Z.; Zeng, M.; Chen, J. Impact of Matrix Species and Mass Spectrometry on Matrix Effects in Multi-Residue Pesticide Analysis Based on QuEChERS-LC-MS. *Foods* **2023**, *12* (6), 1226.

(213) Koloka, O.; Boti, V.; Albanis, T.; Konstantinou, I. Accurate Determination of Pesticide Residues in Milk by Sonication-QuEChERS Extraction and LC-LTQ/Orbitrap Mass Spectrometry. *Separations* **2023**, *10* (3), 146.

(214) Stoyke, M.; Becker, R.; Brockmeyer, J.; Jira, W.; Popping, B.; Uhlig, S.; Wittke, S. German Government Official Methods Board Points the Way Forward: Launch of a New Working Group for Mass Spectrometry for Protein Analysis to Detect Food Fraud and Food Allergens. *Journal of AOAC INTERNATIONAL* **2019**, *102* (5), 1280–1285.

(215) Dou, X.; Zhang, L.; Yang, R.; Wang, X.; Yu, L.; Yue, X.; Ma, F.; Mao, J.; Wang, X.; Zhang, W.; Li, P. Mass Spectrometry in Food Authentication and Origin Traceability. *Mass Spectrom Rev.* **2023**, *42*, No. e21779.

(216) Kosek, V.; Uttl, L.; Jíru, M.; Black, C.; Chevallier, O.; Tomaniová, M.; Elliott, C. T.; Hajišlová, J. Ambient Mass Spectrometry Based on REIMS for the Rapid Detection of Adulteration of Minced Meats by the Use of a Range of Additives. *Food Control* **2019**, *104*, 50–56.

(217) Piras, C.; Hale, O. J.; Reynolds, C. K.; Jones, A. K.; Taylor, N.; Morris, M.; Cramer, R. Speciation and Milk Adulteration Analysis by Rapid Ambient Liquid MALDI Mass Spectrometry Profiling Using Machine Learning. *Sci. Rep* **2021**, *11* (1), 3305.

(218) Nichani, K.; Uhlig, S.; Colson, B.; Hettwer, K.; Simon, K.; Bönick, J.; Uhlig, C.; Kemmlin, S.; Stoyke, M.; Gowik, P.; Huschek, G.; Rawel, H. M. Development of Non-Targeted Mass Spectrometry Method for Distinguishing Spelt and Wheat. *Foods* **2023**, *12* (1), 141.

(219) Ding, B.; Tao, Y.; Xie, J.; Zeng, G.; Huang, H. Traceability Evaluation of Wild and Cultivated *Cordyceps Sinensis* by Elemental Analysis and GasBench II Coupled to Stable Isotope Ratio Mass Spectrometry. *Food Anal. Methods* **2023**, *16* (3), 515–524.

(220) Agriopoulou, S.; Stamatelopoulou, E.; Varzakas, T. Advances in Analysis and Detection of Major Mycotoxins in Foods. *Foods* **2020**, *9* (4), 518.

(221) Liew, W.-P.-P.; Mohd-Redzwan, S. Mycotoxin: Its Impact on Gut Health and Microbiota. *Frontiers in Cellular and Infection Microbiology* **2018**, *8*. DOI: 10.3389/fcimb.2018.00060

(222) Turner, P. C.; Snyder, J. A. Development and Limitations of Exposure Biomarkers to Dietary Contaminants Mycotoxins. *Toxins (Basel)* **2021**, *13* (5), 314.

(223) Donnelly, R.; Elliott, C.; Zhang, G.; Baker, B.; Meneely, J. Understanding Current Methods for Sampling of Aflatoxins in Corn and to Generate a Best Practice Framework. *Toxins* **2022**, *14* (12), 819.

(224) Scheerlinck, E.; Dhaenens, M.; Van Soom, A.; Peelman, L.; De Sutter, P.; Van Steendam, K.; Deforce, D. Minimizing Technical Variation during Sample Preparation Prior to Label-Free Quantitative Mass Spectrometry. *Anal. Biochem.* **2015**, *490*, 14–19.

(225) Piehowski, P. D.; Petyuk, V. A.; Orton, D. J.; Xie, F.; Ramirez-Restrepo, M.; Engel, A.; Lieberman, A. P.; Albin, R. L.; Camp, D. G.; Smith, R. D.; Myers, A. J.; et al. Sources of Technical Variability in Quantitative LC-MS Proteomics: Human Brain Tissue Sample Analysis. *J. Proteome Res.* **2013**, *12* (5), 2128–2137.

(226) Loi, M.; Logrieco, A. F.; Pusztahelyi, T.; Leiter, É.; Hornok, L.; Pócsi, I. Advanced Mycotoxin Control and Decontamination Techniques in View of an Increased Aflatoxin Risk in Europe Due to Climate Change. *Frontiers in Microbiology* **2023**, *13*. DOI: 10.3389/fmicb.2022.1085891

(227) Hansen, H. C. B.; Hilscherova, K.; Bucheli, T. D. Natural Toxins: Environmental Contaminants Calling for Attention. *Environmental Sciences Europe* **2021**, *33* (1), 112.

(228) FSSAI. *MANUAL OF METHODS OF ANALYSIS OF FOODS-Mycotoxins*; Food Safety and Standards Authority of India: India, 2016; p 64. [https://fssai.gov.in/upload/advisories/2018/02/Sa93ebeda71cManual_Mycotoxins_25_05_2016\(1\).pdf](https://fssai.gov.in/upload/advisories/2018/02/Sa93ebeda71cManual_Mycotoxins_25_05_2016(1).pdf).

(229) Cinar, A.; Onbasi, E.; Cinar, A.; Onbasi, E. Mycotoxins: The Hidden Danger in Foods. In *Mycotoxins and Food Safety*; IntechOpen, 2019. DOI: 10.5772/intechopen.89001.

(230) Iqbal, S. Z. Mycotoxins in Food, Recent Development in Food Analysis and Future Challenges; a Review. *Current Opinion in Food Science* **2021**, *42*, 237–247.

(231) Tittlemier, S. a.; Cramer, B.; Dall'Asta, C.; Iha, M. h.; Lattanzio, V. m. t.; Maragos, C.; Solfrizzo, M.; Stranska, M.; Stroka, J.; Sumarah, M. Developments in Mycotoxin Analysis: An Update for 2018–19. *World Mycotoxin Journal* **2020**, *13* (1), 3–24.

(232) Renaud, J. B.; Miller, J. D.; Sumarah, M. W. Mycotoxin Testing Paradigm: Challenges and Opportunities for the Future. *J. AOAC Int.* **2019**, *102* (6), 1681–1688.

(233) Lattanzio, V. M. T.; Solfrizzo, M.; Powers, S.; Visconti, A. Simultaneous Determination of Aflatoxins, Ochratoxin A and Fusarium Toxins in Maize by Liquid Chromatography/Tandem Mass Spectrometry after Multitoxin Immunoaffinity Cleanup. *Rapid Commun. Mass Spectrom.* **2007**, *21* (20), 3253–3261.

(234) Cheng, H.-L.; Wang, F.-L.; Zhao, Y.-G.; Zhang, Y.; Jin, M.-C.; Zhu, Y. Simultaneous Determination of Fifteen Toxic Alkaloids in Meat Dishes and Vegetable Dishes Using Double Layer Pipette Tip Magnetic Dispersive Solid Phase Extraction Followed by UFLC-MS/MS. *Anal. Methods* **2018**, *10* (10), 1151–1162.

(235) Wang, R.; Guo, L.; Wang, P.; Su, X.; Song, Z.; Lin, G.; Zhu, R. [Simultaneous determination of 37 mycotoxins in grain and animal feed by impurity adsorption purification coupled with ultra-performance liquid chromatography-tandem mass spectrometry]. *Se Pu* **2020**, *38* (7), 817–825.

(236) Nualkaw, K.; Poapolathep, S.; Zhang, Z.; Zhang, Q.; Giorgi, M.; Li, P.; Logrieco, A. F.; Poapolathep, A. Simultaneous Determination of Multiple Mycotoxins in Swine, Poultry and Dairy Feeds Using Ultra High Performance Liquid Chromatography-Tandem Mass Spectrometry. *Toxins* **2020**, *12* (4), 253.

(237) Albero, B.; Fernández-Cruz, M. L.; Pérez, R. A. Simultaneous Determination of 15 Mycotoxins in Aquaculture Feed by Liquid Chromatography-Tandem Mass Spectrometry. *Toxins* **2022**, *14* (5), 316.

(238) Berthiller, F.; Crews, C.; Dall'Asta, C.; Saeger, S. D.; Haesaert, G.; Karlovsky, P.; Oswald, I. P.; Seefelder, W.; Speijers, G.; Stroka, J. Masked Mycotoxins: A Review. *Mol. Nutr Food Res.* **2013**, *57* (1), 165–186.

(239) Fiby, I.; Sopol, M. M.; Michlmayr, H.; Adam, G.; Berthiller, F. Development and Validation of an LC-MS/MS Based Method for the Determination of Deoxynivalenol and Its Modified Forms in Maize. *Toxins* **2021**, *13* (9), 600.

(240) Zhang, Y.; Chen, T.; Chen, D.; Liang, W.; Lu, X.; Zhao, C.; Xu, G. Suspect and Nontarget Screening of Mycotoxins and Their Modified Forms in Wheat Products Based on Ultrahigh-Performance Liquid Chromatography-High Resolution Mass Spectrometry. *Journal of Chromatography A* **2023**, *1708*, 464370.

(241) Pascari, X.; Weigel, S.; Marin, S.; Sanchis, V.; Maul, R. Detection and Quantification of Zearalenone and Its Modified Forms

in Enzymatically Treated Oat and Wheat Flour. *J. Food Sci. Technol.* **2023**, *60* (4), 1367–1375.

(242) EMR. *Global Mycotoxin Testing Market Report and Forecast 2023–2028*; EMR, 2022. <https://www.expertmarketresearch.com/reports/mycotoxin-testing-market> (accessed 07-03-2023).

(243) BaySpec Inc. *Continuity™ Portable Mass Spectrometer*; BaySpec, 2023. <https://www.bayspec.com/products/high-sensitivity-portable-mass-spectrometer/> (accessed 07-03-2023).

(244) Nolan, P.; Auer, S.; Spehar, A.; Oplatowska-Stachowiak, M.; Campbell, K. Evaluation of Mass Sensitive Micro-Array Biosensors for Their Feasibility in Multiplex Detection of Low Molecular Weight Toxins Using Mycotoxins as Model Compounds. *Talanta* **2021**, *222*, 121521.